

SI CHROMATIN AND DEVELOPMENT

Meiotic recombination hotspots – a comparative view

Kyuha Choi and Ian R. Henderson*

Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EA, UK

Received 13 March 2015; revised 15 April 2015; accepted 17 April 2015.

*For correspondence (e-mail irh25@cam.ac.uk).

SUMMARY

During meiosis homologous chromosomes pair and undergo reciprocal genetic exchange, termed crossover. Meiotic recombination has a profound effect on patterns of genetic variation and is an important tool during crop breeding. Crossovers initiate from programmed DNA double-stranded breaks that are processed to form single-stranded DNA, which can invade a homologous chromosome. Strand invasion events mature into double Holliday junctions that can be resolved as crossovers. Extensive variation in the frequency of meiotic recombination occurs along chromosomes and is typically focused in narrow hotspots, observed both at the level of DNA breaks and final crossovers. We review methodologies to profile hotspots at different steps of the meiotic recombination pathway that have been used in different eukaryote species. We then discuss what these studies have revealed concerning specification of hotspot locations and activity and the contributions of both genetic and epigenetic factors. Understanding hotspots is important for interpreting patterns of genetic variation in populations and how eukaryotic genomes evolve. In addition, manipulation of hotspots will allow us to accelerate crop breeding, where meiotic recombination distributions can be limiting.

Keywords: meiosis, recombination, hotspots, chromatin, DNA motifs.

INTRODUCTION

Meiosis is widely conserved in sexually reproducing eukaryotes, during which a single round of DNA replication is coupled with two rounds of chromosome segregation (Villeneuve and Hillers, 2001). Prior to the first meiotic division homologous chromosomes pair and undergo programmed recombination, which can result in reciprocal genetic exchange, termed crossover (Hunt Morgan, 1916). In addition to generating new combinations of sequence diversity, an obligate crossover is required to physically connect homologs and ensure balanced chromosome segregation at meiosis I (Page and Hawley, 2003). The frequency of meiotic recombination is highly non-random along chromosomes, and typically occurs within narrow regions termed hotspots (Lichten and Goldman, 1995; Kapppi *et al.*, 2004; de Massy, 2013; Lam and Keeney, 2014; Mercier *et al.*, 2014). In addition, large (megabase) chromosomal regions can be suppressed for recombination, for example the heterochromatic regions around centromeres (Copenhaver *et al.*, 1999; The Tomato Genome Consortium, 2012; Mayer *et al.*, 2012; Choulet *et al.*, 2014). Despite such regions showing low recombination rates in crop genomes, they frequently contain variation in agriculturally

important genes, for example those controlling disease resistance or stress tolerance. Therefore, greater understanding of the control of recombination distributions has the potential to accelerate crop breeding and allow use of the available variation more effectively.

Meiotic recombination is initiated by DNA double-stranded breaks (DSBs) generated by the SPO11 endonuclease (Bergerat *et al.*, 1997; Keeney *et al.*, 1997) (Figure 1a–c). DSB ends then undergo 5' to 3' resection to generate 3'-single-stranded DNA (ssDNA) that is bound by the DMC1 and RAD51 recombinases (Figure 1d,e) (Sun *et al.*, 1991; Bishop *et al.*, 1992; Shinohara *et al.*, 1992). The resulting nucleoprotein filament then performs strand invasion of a homologous chromosome (Kurzbaue *et al.*, 2012; Da Ines *et al.*, 2013). These inter-homolog strand invasion events are further processed to form a double Holliday junction (dHJ) that can be resolved as a crossover (Figure 1f–h) (Szostak *et al.*, 1983; Schwacha and Kleckner, 1995). A larger number of DSBs initially form then mature into final crossover molecules. For example in *Arabidopsis* approximately 200 DSB foci are observed, based on RAD51 and DMC1 foci, whereas only approximately 10 crossovers




Figure 1. Methodologies to profile meiotic recombination hotspots.

(a) Meiotic recombination is initiated by DNA double-stranded breaks (DSBs) generated by SPO11 endonuclease dimers.

(b) SPO11 molecules become covalently bound to DSB target site 5' ends. Upstream and downstream cleavage by the MRX complex releases SPO11-oligonucleotide complexes.

(c) SPO11 can be immunopurified and the bound oligonucleotides analysed to generate maps of meiotic DSBs.

(d) Following DSB formation resection occurs to generate single-stranded DNA (ssDNA) that is bound by the RAD51 and DMC1 recombinases.

(e) RAD51 or DMC1 can be immunopurified and bound ssDNA used to analyse DSB sites. Alternatively, the ssDNA can be directly purified using an affinity resin.

(f) Resected ssDNA invades a homologous chromosome (red) to form a displacement loop.

(g) DNA synthesis and ligation generate a double Holliday junction.

(h) Double Holliday junctions can be resolved as crossover molecules.

(i) Crossover hotspots can be analysed using pollen-typing with allele-specific oligonucleotides (ASOs) that anneal to polymorphic sites between the homologs. This situation allows recombinant (crossover) and non-recombinant (parental) molecules to be specifically amplified.

are detected per meiosis (Copenhaver *et al.*, 1998; Chelysheva *et al.*, 2010; Giraut *et al.*, 2011; Ferdous *et al.*, 2012; Salomé *et al.*, 2012). Most DSBs are thought to enter the strand invasion pathway, but are then resolved as non-crossovers via synthesis-dependent strand annealing (SDSA), without exchange of flanking genetic markers (Allers and Lichten, 2001; McMahon *et al.*, 2007). As this mode of repair involves template-driven DNA synthesis post-invasion, it can result in gene conversion over heterozygous sites, which can be detected as 3:1 inheritance between the meiotic products (Sun *et al.*, 2012; Wijnker *et al.*, 2013).

The majority (approximately 85%) of Arabidopsis crossovers are interfering, meaning they are distributed further apart than expected if they were independent, and are generated by the ZMM pathway (*MSH4*, *MSH5*, *MER3*, *HEI10*, *ZIP4*, *SHOC1*, *PTD*) (Higgins *et al.*, 2004, 2008; Chen *et al.*, 2005; Mercier *et al.*, 2005; Wijeratne *et al.*, 2006; Macaisne *et al.*, 2008; Chelysheva *et al.*, 2012). The ZMM pathway is best characterized in budding yeast in which it acts to stabilize inter-homolog strand invasion intermediates and promotes formation of dHJs and their resolution as cross-

overs (Lynn *et al.*, 2007). The remaining non-ZMM crossovers in Arabidopsis are generated by a non-interfering pathway, that includes *MUS81* (Berchowitz *et al.*, 2007; Higgins, Buckling, *et al.*, 2008). *FANCM*, *MHF1* and *MHF2* are required for non-crossover formation in Arabidopsis, and in their absence DSBs that would normally become non-crossovers are repaired as non-interfering crossovers, leading to an approximately 3-fold increase in total crossover frequency (Crismani *et al.*, 2012; Knoll *et al.*, 2012; Girard *et al.*, 2014). These non-interfering crossovers are assumed to be generated via the *MUS81* pathway, though a formal test of this model is precluded due to *fancm mus81* lethality (Crismani *et al.*, 2012). More recently *TOPOISOMERASE3 α* and the *RECQ4A* and *RECQ4B* helicases have been identified as additional anti-crossover factors in Arabidopsis (Séguéla-Arnaud *et al.*, 2015). As a consequence of these recombination pathways one or a small number of crossovers form per chromosome, although the positions of crossovers observed between independent meioses are highly variable.

Measurement of DSBs and crossovers in multiple species has demonstrated concentration in hotspots, which

are typically 1–10 kb in width and show significantly higher recombination activity than surrounding regions (Lichten and Goldman, 1995; Kauppi *et al.*, 2004; de Massy, 2013; Lam and Keeney, 2014; Mercier *et al.*, 2014). In this review we examine the methodologies used in different eukaryotic species to profile meiotic break and crossover hotspots (Table 1). We then discuss what these studies have revealed concerning the genetic and epigenetic factors that contribute to variation in recombination frequency along eukaryotic chromosomes and the formation of hotspots.

DETECTING AND MEASURING HOTSPOTS

DSB hotspots – mapping breaks and ssDNA

Meiotic recombination initiates from DNA DSBs, which can be directly measured to reveal hotspots. A major advantage to the study of meiosis in budding yeast *Saccharomyces cerevisiae* is the ability to synchronize large numbers of cells to enter meiosis simultaneously, which facilitates biochemical analysis of meiotic recombination. Due to the effect of DSBs on DNA molecule size, gel electrophoresis of staged meiotic DNA can be used to analyse DSBs in yeast (Nicolas *et al.*, 1989; Baudat and Nicolas, 1997; Garcia *et al.*, 2015). This is a versatile method that allows DSB variation to be measured on scales from whole chromosomes to single nucleotides, according to electrophoresis conditions and probe choice (Nicolas *et al.*, 1989; Baudat and Nicolas, 1997; Garcia *et al.*, 2015). The detection of DSBs can be increased by using DNA processing mutants such as *rad50S* and *sae2D*, which are deficient in endonucleolytic release of SPO11 from DSBs and accumulate unrepaired breaks (Buhler *et al.*, 2007). However, it is important to note that mapping DSBs in *rad50s* and *sae2D* mutants may lead to biased detection of early forming DSBs over late DSBs, which account for a substantial fraction of breaks in *S. cerevisiae* (Buhler *et al.*, 2007; Joshi *et al.*, 2015). Although the application of gel electrophoresis assays to larger genomes is challenging, a modified Southern blotting method using terminal transferase and nested PCR was able to detect DSBs at the *H2-Ea* hotspot in mouse testicular germ cells (Qin *et al.*, 2004).

A sensitive method to detect meiotic DSBs relies on the catalytic mechanism of the SPO11 endonuclease (Neale *et al.*, 2005). SPO11 is related to topoisomerases and becomes covalently bound to 5'-target sites at a catalytic tyrosine residue (Bergerat *et al.*, 1997; Keeney *et al.*, 1997; Neale *et al.*, 2005). Subsequent upstream or downstream cleavage by the MRX nuclease complex liberates SPO11 bound to oligonucleotides approximately 20–100 nt in length (Figure 1b) (Neale *et al.*, 2005; Garcia *et al.*, 2011; Lange *et al.*, 2011; Pan *et al.*, 2011). Immunoprecipitation of SPO11 and purification of the bound oligonucleotides can be used to analyse DSB target sites, using end-labelling and gel electrophoresis, or generation of sequencing libraries (Figure 1c) (Neale *et al.*, 2005; Lange *et al.*, 2011; Pan *et al.*, 2011). These approaches have been performed successfully in budding yeast, fission yeast and mice (Lange *et al.*, 2011; Pan *et al.*, 2011; Fowler *et al.*, 2014). This technique has also been repeated in mouse *atm* kinase signaling mutants, which accumulate higher DSB levels (Lange *et al.*, 2011). In budding yeast Spo11 functions with a group of accessory proteins, including Rec114, Mer2 and Mei4 (Panizza *et al.*, 2011). These proteins have been analysed by chromatin immunoprecipitation (ChIP) and shown to associate with cohesin-rich regions (Panizza *et al.*, 2011), which may reflect a mechanism to tether DSBs formed on chromatin loops to repair sites at the meiotic chromosome axis (Kleckner *et al.*, 2004). Alternatively, the locus where breaks will form is first tethered to the axis, where SPO11 is present, and then a DSB forms (Kleckner *et al.*, 2004; Sommermeyer *et al.*, 2013).

Following DSB resection, meiotic ssDNA can be directly purified and analysed using an affinity resin, which can also be combined with recombination/processing mutants that accumulate unrepaired breaks (Figure 1e) (Buhler *et al.*, 2007, 2009). As ssDNA is bound by the DMC1 and RAD51 recombinases, ChIP of these factors has been used to generate high-resolution recombination maps in yeast, mouse and humans, using an approach termed ssDNA-sequencing (SSDS) (Figure 1e) (Smagulova *et al.*, 2011; Brick *et al.*, 2012; Khil *et al.*, 2012; He *et al.*, 2013; Pratto *et al.*, 2014). Similar methods have also been developed in

Table 1 Methodologies used to detect meiotic recombination hotspots throughout eukaryotes

Recombination step	Technique	Fungi	Plants	Animals
DNA double strand breaks	Gel electrophoresis & Southern blotting	o		o
DNA double strand breaks	RPA & DMC1 ChIP-seq (SSDS)	o	o	o
DNA double strand breaks	ssDNA purification	o		
DNA double strand breaks	SPO11 oligonucleotide sequencing	o		o
ZMM pathway	Zip3 chromatin immunoprecipitation	o		
Crossovers/non-crossovers	Tetrad analysis	o	o	o
Crossovers/non-crossovers	Classical genetic mapping	o	o	o
Crossovers/non-crossovers	Gamete/spore genome sequencing	o	o	o
Crossovers/non-crossovers	Sperm-typing/pollen-typing		o	o
Crossovers/non-crossovers	LD-based analyses	o	o	o

maize (He *et al.*, 2013). It will be important to apply these or related methods in plants to fully understand genomic distributions of meiotic DSBs. A variety of *Arabidopsis* meiotic mutants are available that may facilitate the study of DSBs, including those altering break processing (*sae2/com1*) (Uanschou *et al.*, 2007), DNA damage kinase signaling (*atm, atr*) (Garcia *et al.*, 2003; Roitinger *et al.*, 2015) and chromatin modification (*arp6, met1, ddm1*) (Colomé-Tatché *et al.*, 2012; Melamed-Bessudo and Levy, 2012; Mirouze *et al.*, 2012; Yelina *et al.*, 2012; Choi *et al.*, 2013).

Crossover hotspots – direct mapping

A classical method to detect crossovers is via analysis of co-inheritance of linked heterozygous markers through meiosis (Hunt Morgan, 1916), in which change of linkage phase between markers indicates the occurrence of a crossover. Meiotic tetrad analysis is a powerful genetic technique, where all four daughter cells from a single meiosis are analysed (Lichten, 2014). In addition to distinguishing between two, three and four strand crossover events, tetrad analysis also allows measurement of gene conversion events via detection of 3:1 inheritance between sister gametes (Nicolas *et al.*, 1989; Berchowitz and Copenhaver, 2008; Martini *et al.*, 2011; Sun *et al.*, 2012). Recently tetrad analysis has been extended to both mice and plants (Berchowitz and Copenhaver, 2008; Cole *et al.*, 2014). In *Arabidopsis* the *quartet1 (qrt1)* mutant is altered in pollen wall biogenesis, such that the four products of male meiosis remain physically attached (Francis *et al.*, 2006). Combination of *qrt1* with linked heterozygous transgenes expressing different colours of fluorescent protein provides an elegant visual method to score crossovers (Francis *et al.*, 2007; Berchowitz and Copenhaver, 2008), and an equivalent approach has been developed in budding yeast (Thacker *et al.*, 2011). Alternatively, *qrt1* can be complemented and single pollen grains analysed using flow

cytometry to increase measurement throughput (Yelina *et al.*, 2012, 2013). Single *qrt1* tetrad pollinations can also be used to isolate siblings related through a common meiosis, which are then sequenced to provide insight into genome-wide patterns of crossover and gene conversion (Wijnker *et al.*, 2013; Qi *et al.*, 2014). Equally, large numbers of crossover events can be readily mapped using genotyping or sequencing of F₂ or backcross populations (Salomé *et al.*, 2012; Rowan *et al.*, 2015). New methods that use multiple annealing- and looping-based amplification cycles (MALBEC) have allowed sequencing and identification of crossovers in a single human sperm or oocyte (Lu *et al.*, 2012; Hou *et al.*, 2013). This outcome has also been achieved in maize via sequencing of DNA from isolated single microspores (the male meiotic products) (Li *et al.*, 2015). However, a major limitation to all of these approaches for the study of hotspots is the paucity of crossovers per meiosis. For example, known plant hotspots have genetic distances between approximately 0.1–0.5 cM (Table 2) (Yelina *et al.*, 2012; Choi *et al.*, 2013; Drouaud *et al.*, 2013). Therefore, to analyse 100s of crossovers at a given hotspot it is necessary to screen 100 000s of meioses, which is generally not possible when generating populations of plants.

To isolate large numbers of crossovers at individual hotspots a general approach is to use allele-specific amplification from post-meiotic gamete DNA, termed sperm typing or pollen typing (Tiemann-Boege *et al.*, 2006; Baudat and de Massy, 2007; Cole *et al.*, 2010; Berg *et al.*, 2011; Drouaud and Mézard, 2011; Yelina *et al.*, 2012; Choi *et al.*, 2013; Drouaud *et al.*, 2013). These approaches rely on the generation of individuals that are heterozygous over a hotspot region of interest. Collection of gametes and isolation of DNA means that samples consist of a mixture of non-recombinant (parental) and crossover molecules that are distinguishable by patterns of polymorphisms. Allele-spe-

Table 2 Plant meiotic recombination hotspots

Species	Hotspot	Interval (base-pairs)	cM	cM/Mb	Location	Chromosome cM/Mb	Reference
<i>Arabidopsis thaliana</i>	3b	5746	0.11	20.01	Intergenic	4.77	Choi <i>et al.</i> (2013)
<i>Arabidopsis thaliana</i>	3a	5825	0.21	36.22	Gene 5' and 3' end	4.77	Choi <i>et al.</i> (2013); Yelina <i>et al.</i> (2012)
<i>Arabidopsis thaliana</i>	14a	7283	0.55	75.52	Gene 5' end & intergenic	4.8	Drouaud <i>et al.</i> (2013)
<i>Arabidopsis thaliana</i>	130x	12 488	0.53	42.44	Intergenic	4.8	Drouaud <i>et al.</i> (2013)
<i>Zea mays</i>	a1	1900	0.08	40.00	a1	2.1	Brown and Sundaresan (1991)
<i>Zea mays</i>	a1	710	0.04	59.00	Gene promoter	0.5–1.5	Yao and Schnable (2005)
<i>Zea mays</i>	a1	500	0.005	9.10	Gene 5' end	0.5–1.5	Yao and Schnable (2005)
<i>Zea mays</i>	Yz1	1900	0.06	32.00	Gene 5' end	0.5–1.5	Yao and Schnable (2005)
<i>Zea mays</i>	Yz1	540	0.03	48.00	Gene 3' end	0.5–1.5	Yao and Schnable (2005)
<i>Zea mays</i>	B	620	0.03	52.00	Gene 5' end	nd	Patterson <i>et al.</i> (1995)
<i>Zea mays</i>	bronze	1056	0.051	48.30	Gene 5' end	nd	Fu <i>et al.</i> (2002)
<i>Zea mays</i>	bronze	793	0.033	41.61	Gene 5' end	nd	Fu <i>et al.</i> (2002)
<i>Triticum aestivum</i>	HGA3	23 000	0.20	8.82	Gene 5' end	0.85	Saintenac <i>et al.</i> (2009)

cific primers are designed to anneal to polymorphisms flanking a hotspot and used in different configurations to amplify either parental or crossover molecules (Figure 1i) (Baudat and de Massy, 2009; Kauppi *et al.*, 2009; Cole and Jasin, 2011; Drouaud and Mézard, 2011). Typically, the products of male meiosis (pollen or sperm) are collected due to the ease of isolating the required large numbers of cells (Kauppi *et al.*, 2009; Drouaud and Mézard, 2011), although analysis of female gametes is also possible (de Boer *et al.*, 2013). This analysis is important as sex-specific differences in meiotic recombination are widespread, yet poorly understood (Lenormand and Dutheil, 2005; Giraut *et al.*, 2011; Campbell *et al.*, 2015). Estimating recombination rate within the amplified region is possible by diluting the gamete template DNA in order to titrate parental and crossover molecules. It is possible to then sequence or genotype amplification products from single crossover molecules to identify internal crossover positions to the resolution of individual polymorphisms (Tiemann-Boege *et al.*, 2006; Baudat and de Massy, 2007; Cole *et al.*, 2010; Berg *et al.*, 2011; Drouaud and Mézard, 2011; Yelina *et al.*, 2012; Choi *et al.*, 2013; Drouaud *et al.*, 2013). These methods allow the fine-scale characterization of crossover patterns within single hotspots. A further method to map crossover recombination is via ChIP of associated factors, for example Zip3 within the ZMM pathway has been analysed in budding yeast (Serrentino *et al.*, 2013). In some cases early and late cytogenetic foci are observed with distinct properties (e.g. RNF212), and so it is important to consider whether ChIP approaches will profile different foci classes equally (Reynolds *et al.*, 2013; Qiao *et al.*, 2014).

Crossover hotspots – historical mapping

The signature of crossover can also be detected via analysis of natural genetic polymorphisms, due to the effect recombination has on non-random associations between mutations (Auton and McVean, 2012). Specifically, crossovers cause decay of linkage disequilibrium (LD) between linked polymorphisms, which can be analysed using coalescent theory (Charlesworth and Charlesworth, 2010; Auton and McVean, 2012). For example, consider two linked, independently arising mutations *a* and *b*. The only way an *a–b* haplotype can occur is through recurrent mutation, or via recombination joining *a* and *b* onto the same chromosome (Hudson and Kaplan, 1985; Charlesworth and Charlesworth, 2010; Auton and McVean, 2012). Packages such as LDhat and SequenceLDhot use this principle to analyse SNP patterns and estimate the population-scaled recombination rate $4N_e r$ (where *r* is the per generation recombination rate and N_e is the effective population size) (Fearnhead, 2006; Auton and McVean, 2007). These approaches are powerful as they sample the very large numbers of meioses occurring in the history of the individuals compared. However, there are a number of limitations

to these methods, including that SNPs are influenced by population genetic forces in addition to recombination, including selection, drift and migration (Charlesworth and Charlesworth, 2010; Auton and McVean, 2012). Further caveats associated with these approaches include the potential error introduced by structural variation between individuals, for example insertions, deletions, inversions and translocations. Mis-calling of SNP positions relative to a given reference sequence may under- or over-estimate actual physical distances between variants and thus cause erroneous recombination rate estimates. This may be a particular problem in repetitive regions, where accurately identifying SNPs from short read sequencing data can be problematic. Therefore, it is important to combine historical and experimental mapping of crossover recombination.

GENETIC AND EPIGENETIC CONTROL OF HOTSPOTS

Gene-associated recombination hotspots in plants and fungi

Recombination mapping techniques have revealed much concerning hotspot locations and their control in plant, animal and fungal species (Lichten and Goldman, 1995; Kauppi *et al.*, 2004; de Massy, 2013; Lam and Keeney, 2014; Mercier *et al.*, 2014). Despite the widespread occurrence of hotspots throughout eukaryotes major differences in the relative importance of genetic and epigenetic control exist. In *Arabidopsis* a combination of historical and experimental mapping of crossovers has revealed concentration of recombination at gene promoters and terminators (Figure 2a) (Yelina *et al.*, 2012; Choi *et al.*, 2013; Drouaud *et al.*, 2013). The historical crossover signal in this species is highest immediately downstream of gene transcriptional start sites (TSSs), overlapping the highly positioned +1 nucleosome (Figure 2a) (Choi *et al.*, 2013). Significant but lower peaks are also observed in proximity to gene transcriptional termination sites (TTSs), also within gene open reading frames (Figure 2a) (Choi *et al.*, 2013). Historical crossover hotspots at the start and end of genes were also observed in *Mimulus guttatus* (Hellsten *et al.*, 2013). Direct mapping of crossovers has shown gene-associated hotspots in maize (*Bronze*, *a1*) and wheat (*HGA1*, *HGA3*, *SOS1*) (Table 2) (Dooner, 1986; Xu *et al.*, 1995; Fu *et al.*, 2001; Sainenac *et al.*, 2009; Li *et al.*, 2015), and is concordant with the high recombination rates in gene-rich euchromatin and low rates in repeat-rich heterochromatin observed in multiple plant genomes (Gore *et al.*, 2009; Liu *et al.*, 2009; Wei *et al.*, 2009; The Tomato Genome Consortium, 2012; Mayer *et al.*, 2012; Choulet *et al.*, 2014; Li *et al.*, 2015; Rodgers-Melnick *et al.*, 2015).

Productive and accurate gene transcription by RNA polymerase II is known to rely on both specific DNA sequences and chromatin states at gene promoters (Venters and Pugh, 2009; Deal and Henikoff, 2011). For example, the +1




Figure 2. Plant versus human crossover hotspots.

(a) Crossover frequency (green) is represented at plant gene-associated hotspots. Crossovers are highest in proximity to gene transcriptional start (TSSs) and termination (TTSs) sites. The gene open reading frame is shown by the black arrow. Recombination hotspots at the gene promoter (TSS) are also associated with +1 nucleosomes (red circle) that contain H2A.Z and are modified with H3K4^{me3} and CTT-repeat DNA sequence motifs (Choi *et al.*, 2013; Wijnker *et al.*, 2013).

(b) In humans crossover hotspots occur at intergenic CCN-like motifs (Myers *et al.*, 2008), which are bound by PRDM9 zinc finger domains. PRDM9 also contains a histone methyltransferase SET domain, which catalyses H3K4^{me3} (red circles) and causes high recombination rates.

nucleosome shows the H3K4^{me3} modification and the presence of the histone variant H2A.Z, both of which are important for regulation of transcription (Zhang *et al.*, 2009; Coleman-Derr and Zilberman, 2012). The deposition of H2A.Z has also been shown to be required for normal levels of meiotic recombination at multiple physical scales, via analysis of the *arp6* mutation (Choi *et al.*, 2013). ARP6 is a component of the SWR1 chromatin remodeling complex that deposits H2A.Z into chromatin (Choi *et al.*, 2005). RAD51 and DMC1 foci were also significantly reduced in *arp6*, which indicates that DSB frequency may be lower, in addition to final crossover numbers (Choi *et al.*, 2013). It is also important to note that *arp6* plays an important role in the transcription of meiotic genes, including *DMC1*, which may contribute to mutant phenotypes (Qin *et al.*, 2014).

The presence of hotspots at plant gene promoters is similar to the occurrence of DSB hotspots at gene promoters in budding yeast (Nicolas *et al.*, 1989; Wu and Lichten, 1994; Lichten and Goldman, 1995; Berchowitz *et al.*, 2009; Pan *et al.*, 2011). In budding yeast nucleosome occupancy is a major determinant of recombination and high levels of DSBs are observed in nucleosome-free regions upstream of TSSs (Nicolas *et al.*, 1989; Wu and Lichten, 1994; Lichten and Goldman, 1995; Berchowitz *et al.*, 2009; Pan *et al.*,

2011). The H3K4^{me3} chromatin modification plays an important role in promoting hotspot recombination in budding yeast (Borde *et al.*, 2009; Tischfield and Keeney, 2012). Mutations to the *SET1* histone H3K4 methyltransferase cause severely reduced DSB frequency at the majority of hotspots, although a smaller number showed increased break frequency (Borde *et al.*, 2009). The Spp1 subunit of the COMPASS complex was found to interact simultaneously with both H3K4^{me3} and a component of the meiotic axis (Mer2), revealing how epigenetic marks can function to promote tethering of DSBs to axis-associated repair sites (Acquaviva *et al.*, 2013; Sommermeyer *et al.*, 2013). As H3K4^{me3} is also enriched at plant gene promoter hotspots it will be important to test the extent to which this mechanism is conserved. Importantly, hotspot distributions can be evolutionarily diverse even between more closely related species. For example, mapping Spo11-oligos in *Schizosaccharomyces pombe* has revealed distinct patterns relative to budding yeast (Cromie *et al.*, 2007; Hyppa and Smith, 2010; Fowler *et al.*, 2014). Specifically, DSB hotspots are broader, observed in intergenic regions and show a weaker correlation with nucleosome occupancy (Cromie *et al.*, 2007; Hyppa and Smith, 2010; Fowler *et al.*, 2014). Interestingly, fission yeast hotspots also show a bias for inter-sister repair, leading to the phenomenon of crossover invariance (Young *et al.*, 2002; Hyppa and Smith, 2010). Crossover invariance is defined by the observations that in fission yeast large variations in DSB frequency are observed per physical distance, yet crossover frequency is not observed to show matching variation (Hyppa and Smith, 2010). Together this demonstrates the importance of studying hotspots and meiotic recombination in multiple species.

Although chromatin plays a major role in directing recombination at plant hotspots an involvement for specific DNA sequence motifs has also been observed (Choi *et al.*, 2013; Wijnker *et al.*, 2013). An association was detected between crossover hotspots and poly dA:dT sequences located upstream of gene TSS sites, which are known to disfavor nucleosome occupancy (Iyer and Struhl, 1995; Segal and Widom, 2009). One possibility is that A-rich sequences cause a greater degree of nucleosome exclusion that facilitates accessibility of the recombination machinery, including during DSB formation by SPO11 (Nicolas *et al.*, 1989; Wu and Lichten, 1994; Lichten and Goldman, 1995; Berchowitz *et al.*, 2009; Pan *et al.*, 2011). In addition to A-rich sequences, a class of CTT-repeat sequence motifs was detected as hotspot-enriched and located immediately downstream of TSSs, overlapping the +1 H2A.Z-containing nucleosome (Figure 2a) (Choi *et al.*, 2013; Wijnker *et al.*, 2013). These degenerate motifs are of unknown function but it is possible that they also contribute to organization of promoter chromatin in plants with consequences for meiotic recombination. Alternatively the CTT motifs may be specifically recognized by a component

of the recombination machinery, in a manner similar to PRDM9 in animals (Jeffreys *et al.*, 2001; Tiemann-Boege *et al.*, 2006; Baudat and de Massy, 2007; Berg *et al.*, 2010; Cole *et al.*, 2010).

PRDM9-dependent hotspots in animals

Extensive direct mapping of crossovers using sperm typing approaches in humans and mice has revealed the presence of punctate hotspots (Jeffreys *et al.*, 2001; Tiemann-Boege *et al.*, 2006; Baudat and de Massy, 2007; Berg *et al.*, 2010; Cole *et al.*, 2010). This finding is supported by LD-based analysis that revealed thousands of hotspots throughout the human genome (Crawford *et al.*, 2004; McVean *et al.*, 2004; Myers *et al.*, 2005; Khil and Camerini-Otero, 2010). Analysis of human hotspots identified a family of degenerate C-rich DNA sequence motifs associated with hotspots (Myers *et al.*, 2008). Subsequently this motif family was found to reflect binding sites for the PRDM9 protein, which contains both zinc fingers and a SET histone methyltransferase domain (Figure 2b) (Baudat *et al.*, 2010; Myers *et al.*, 2010; Parvanov *et al.*, 2010; Grey *et al.*, 2011). PRDM9 has been shown to direct meiosis-specific H3K4^{me3} that reorganizes nucleosomes around target motifs and drives DSB formation (Figure 2b) (Mihola *et al.*, 2009; Brick *et al.*, 2012; Baker *et al.*, 2014; Pratto *et al.*, 2014). Intriguingly, PRDM9 is an extremely fast-evolving protein, which is reflected in distinctive hotspot locations between human populations and mammalian species (Oliver *et al.*, 2009; Myers *et al.*, 2010; Hinch *et al.*, 2011; Auton *et al.*, 2012). The action of PRDM9 tends to place hotspots in intergenic regions located away from genes (Figure 2b). PRDM9 has mutated to non-functionality in some species, including the canine lineage, where it is associated with reversion of recombination to promoters (Oliver *et al.*, 2009; Auton *et al.*, 2013). This has also been directly demonstrated in *prdm9* mutant mice, where hotspots were found to revert to promoter locations via SSDS mapping (Brick *et al.*, 2012). Clear orthologs of PRDM9 have so far not been identified in plants or fungi. Together these results are consistent with gene promoters representing an ancestral mode of hotspot designation and PRDM9 representing a derived mechanism within animals. These findings again clearly demonstrate the importance of studying meiotic recombination hotspots in multiple eukaryotic lineages.

FUTURE DIRECTIONS

The occurrence of meiotic recombination hotspots is widespread within eukaryotes, but surprisingly the mechanisms that specify their locations and activity are varied. It will be important to study these phenomena in further diverse species to assess evolutionary changes in the relative roles of genetic and epigenetic information in hotspot control. Further work is required to compare hotspots measured early (e.g. DSBs) and late (e.g. crossovers) in the meiotic

recombination pathway to understand their inter-relationships, particularly in relation to homeostatic mechanisms such as crossover interference. Equally, how different levels of meiotic chromosome organization interact to control hotspots and broad scale rates of recombination will be important, including interactions between chromatin and the meiotic axis/synaptonemal complex. The biological significance of hotspots will be important to further explore. For example, are specific types of genes enriched for hotspots and does recombination play an adaptive role at such genes? Finally, a mechanistic understanding of hotspots will allow us to manipulate this process within plant genomes and facilitate crop breeding and improvement.

REFERENCES

- Acquaviva, L., Székvölgyi, L., Dichtl, B., Dichtl, B.S., de La Roche Saint André, C., Nicolas, A. and Géli, V. (2013) The COMPASS subunit Spp1 links histone methylation to initiation of meiotic recombination. *Science*, **339**, 215–218.
- Allers, T. and Lichten, M. (2001) Differential timing and control of noncrossover and crossover recombination during meiosis. *Cell*, **106**, 47–57.
- Auton, A., Fedel-Alon, A., Pfeifer, S. *et al.* (2012) A fine-scale chimpanzee genetic map from population sequencing. *Science*, **336**, 193–198.
- Auton, A. and McVean, G. (2012) Estimating recombination rates from genetic variation in humans. *Methods Mol. Biol.* **856**, 217–237.
- Auton, A. and McVean, G. (2007) Recombination rate estimation in the presence of hotspots. *Genome Res.* **17**, 1219–1227.
- Auton, A., Rui Li, Y., Kidd, J. *et al.* (2013) Genetic recombination is targeted towards gene promoter regions in dogs. *PLoS Genet.* **9**, e1003984.
- Baker, C.L., Walker, M., Kajita, S., Petkov, P.M. and Paigen, K. (2014) PRDM9 binding organizes hotspot nucleosomes and limits Holliday junction migration. *Genome Res.* **24**, 724–732.
- Baudat, F., Buard, J., Grey, C., Fedel-Alon, A., Ober, C., Przeworski, M., Coop, G. and de Massy, B. (2010) PRDM9 is a major determinant of meiotic recombination hotspots in humans and mice. *Science*, **327**, 836–840.
- Baudat, F. and de Massy, B. (2007) Cis- and trans-acting elements regulate the mouse *Psbmb9* meiotic recombination hotspot. *PLoS Genet.* **3**, e100.
- Baudat, F. and de Massy, B. (2009) Parallel detection of crossovers and non-crossovers in mouse germ cells. *Methods Mol. Biol.* **557**, 305–322.
- Baudat, F. and Nicolas, A. (1997) Clustering of meiotic double-strand breaks on yeast chromosome III. *Proc. Natl Acad. Sci. USA*, **94**, 5213–5218.
- Berchowitz, L.E. and Copenhaver, G.P. (2008) Fluorescent *Arabidopsis* tetrads: a visual assay for quickly developing large crossover and crossover interference data sets. *Nat. Protoc.* **3**, 41–50.
- Berchowitz, L.E., Francis, K.E., Bey, A.L. and Copenhaver, G.P. (2007) The role of *AtMUS81* in interference-insensitive crossovers in *A. thaliana*. *PLoS Genet.* **3**, 10.
- Berchowitz, L.E., Hanlon, S.E., Lieb, J.D. and Copenhaver, G.P. (2009) A positive but complex association between meiotic double-strand break hotspots and open chromatin in *Saccharomyces cerevisiae*. *Genome Res.* **19**, 2245–2257.
- Berg, I.L., Neumann, R., Lam, K.-W.G., Sarbajna, S., Odenthal-Hesse, L., May, C.A. and Jeffreys, A.J. (2010) PRDM9 variation strongly influences recombination hot-spot activity and meiotic instability in humans. *Nat. Genet.* **42**, 859–863.
- Berg, I.L., Neumann, R., Sarbajna, S., Odenthal-Hesse, L., Butler, N.J. and Jeffreys, A.J. (2011) Variants of the protein PRDM9 differentially regulate a set of human meiotic recombination hotspots highly active in African populations. *Proc. Natl Acad. Sci. USA*, **108**, 12378–12383.
- Bergerat, A., de Massy, B., Gabelle, D., Varoutas, P.C., Nicolas, A. and Forterre, P. (1997) An atypical topoisomerase II from Archaea with implications for meiotic recombination. *Nature*, **386**, 414–417.
- Bishop, D.K., Park, D., Xu, L. and Kleckner, N. (1992) DMC1: a meiosis-specific yeast homolog of *E. coli* recA required for recombination, synaptonemal complex formation, and cell cycle progression. *Cell*, **69**, 439–456.
- de Boer, E., Jasin, M. and Keeney, S. (2013) Analysis of recombinants in female mouse meiosis. *Methods Mol. Biol.* **957**, 19–45.

- Borde, V., Robine, N., Lin, W., Bonfils, S., Géli, V. and Nicolas, A. (2009) Histone H3 lysine 4 trimethylation marks meiotic recombination initiation sites. *EMBO J.* **28**, 99–111.
- Brick, K., Smagulova, F., Khil, P., Camerini-Otero, R.D. and Petukhova, G.V. (2012) Genetic recombination is directed away from functional genomic elements in mice. *Nature*, **485**, 642–645.
- Brown, J. and Sundaesan, V. (1991) A recombination hotspot in the maize A1 intragenic region. *Theor. Appl. Genet.* **81**, 185–188.
- Buhler, C., Borde, V. and Lichten, M. (2007) Mapping meiotic single-strand DNA reveals a new landscape of DNA double-strand breaks in *Saccharomyces cerevisiae*. *PLoS Biol.* **5**, e324.
- Buhler, C., Shroff, R. and Lichten, M. (2009) Genome-wide mapping of meiotic DNA double-strand breaks in *Saccharomyces cerevisiae*. *Methods Mol. Biol.* **557**, 143–164.
- Campbell, C.L., Furlotte, N.A., Eriksson, N., Hinds, D. and Auton, A. (2015) Escape from crossover interference increases with maternal age. *Nat. Commun.* **6**, 6260.
- Charlesworth, B. and Charlesworth, D. (2010) *Elements of Evolutionary Genetics*. Colorado: Roberts and Company Publishers.
- Chelysheva, L., Grandont, L., Vrielynck, N., le Guin, S., Mercier, R. and Grellon, M. (2010) An easy protocol for studying chromatin and recombination protein dynamics during *Arabidopsis thaliana* meiosis: immunodetection of cohesins, histones and MLH1. *Cytogenet. Genome Res.* **129**, 143–153.
- Chelysheva, L., Vezon, D., Chambon, A. et al. (2012) The *Arabidopsis* HEI10 is a new ZMM protein related to Zip3. *PLoS Genet.* **8**, e1002799.
- Chen, C., Zhang, W., Timofejeva, L., Gerardin, Y. and Ma, H. (2005) The *Arabidopsis* ROCK-N-ROLLERS gene encodes a homolog of the yeast ATP-dependent DNA helicase MER3 and is required for normal meiotic crossover formation. *Plant J.* **43**, 321–334.
- Choi, K., Kim, S., Kim, S.Y. et al. (2005) SUPPRESSOR OF FRIGIDA3 encodes a nuclear ACTIN-RELATED PROTEIN6 required for floral repression in *Arabidopsis*. *Plant Cell*, **17**, 2647–2660.
- Choi, K., Zhao, X., Kelly, K.A. et al. (2013) *Arabidopsis* meiotic crossover hot spots overlap with H2A.Z nucleosomes at gene promoters. *Nat. Genet.* **45**, 1327–1336.
- Choulet, F., Alberti, A., Theil, S. et al. (2014) Structural and functional partitioning of bread wheat chromosome 3B. *Science* **345**, 1249721.
- Cole, F., Baudat, F., Grey, C., Keeney, S., de Massy, B. and Jasin, M. (2014) Mouse tetrad analysis provides insights into recombination mechanisms and hotspot evolutionary dynamics. *Nat. Genet.* **46**, 1072–1080.
- Cole, F. and Jasin, M. (2011) *DNA Recombination*. (Tsubouchi, H. ed.). Totowa, NJ: Humana Press.
- Cole, F., Keeney, S. and Jasin, M. (2010) Comprehensive, fine-scale dissection of homologous recombination outcomes at a hot spot in mouse meiosis. *Mol. Cell*, **39**, 700–710.
- Coleman-Derr, D. and Zilberman, D. (2012) Deposition of histone variant H2A.Z within gene bodies regulates responsive genes. *PLoS Genet.* **8**, e1002988.
- Colomé-Tatché, M., Cortijo, S., Wardenaar, R. et al. (2012) Features of the *Arabidopsis* recombination landscape resulting from the combined loss of sequence variation and DNA methylation. *Proc. Natl Acad. Sci. USA*, **109**, 16240–16245.
- Copenhaver, G.P., Browne, W.E. and Preuss, D. (1998) Assaying genome-wide recombination and centromere functions with *Arabidopsis* tetrads. *Proc. Natl Acad. Sci. USA*, **95**, 247–252.
- Copenhaver, G.P., Nickel, K., Kuromori, T. et al. (1999) Genetic definition and sequence analysis of *Arabidopsis* centromeres. *Science*, **286**, 2468–2474.
- Crawford, D.C., Bhargale, T., Li, N., Hellenthal, G., Rieder, M.J., Nickerson, D.A. and Stephens, M. (2004) Evidence for substantial fine-scale variation in recombination rates across the human genome. *Nat. Genet.* **36**, 700–706.
- Crismani, W., Girard, C., Froger, N., Pradillo, M., Santos, J.L., Chelysheva, L., Copenhaver, G.P., Horlow, C. and Mercier, R. (2012) FANCM limits meiotic crossovers. *Science*, **336**, 1588–1590.
- Cromie, G.A., Hyppa, R.W., Cam, H.P., Farah, J.A., Grewal, S.I.S. and Smith, G.R. (2007) A discrete class of intergenic DNA dictates meiotic DNA break hotspots in fission yeast. *PLoS Genet.* **3**, e141.
- Deal, R.B. and Henikoff, S. (2011) Histone variants and modifications in plant gene regulation. *Curr. Opin. Plant Biol.* **14**, 116–122.
- Dooner, H.K. (1986) Genetic fine structure of the BRONZE locus in maize. *Genetics*, **113**, 1021–1036.
- Drouaud, J., Khademian, H., Giraut, L., Zanni, V., Bellalou, S., Henderson, I.R., Falque, M. and Mézard, C. (2013) Contrasted patterns of crossover and non-crossover at *Arabidopsis thaliana* meiotic recombination hotspots. *PLoS Genet.* **9**, e1003922.
- Drouaud, J. and Mézard, C. (2011) Characterization of meiotic crossovers in pollen from *Arabidopsis thaliana*. *Methods Mol. Biol.* **745**, 223–249.
- Fearnhead, P. (2006) SequenceLDhot: detecting recombination hotspots. *Bioinformatics*, **22**, 3061–3066.
- Ferdous, M., Higgins, J.D., Osman, K. et al. (2012) Inter-homolog crossing-over and synapsis in *Arabidopsis* meiosis are dependent on the chromosome axis protein AtASY3. *PLoS Genet.* **8**, e1002507.
- Fowler, K.R., Sasaki, M., Milman, N., Keeney, S. and Smith, G.R. (2014) Evolutionarily diverse determinants of meiotic DNA break and recombination landscapes across the genome. *Genome Res.* **24**, 1650–1664.
- Francis, K.E., Lam, S.Y. and Copenhaver, G.P. (2006) Separation of *Arabidopsis* pollen tetrads is regulated by QUARTET1, a pectin methylesterase gene. *Plant Physiol.* **142**, 1004–1013.
- Francis, K.E., Lam, S.Y., Harrison, B.D., Bey, A.L., Berchowitz, L.E. and Copenhaver, G.P. (2007) Pollen tetrad-based visual assay for meiotic recombination in *Arabidopsis*. *Proc. Natl Acad. Sci. USA*, **104**, 3913–3918.
- Fu, H., Park, W., Yan, X., Zheng, Z., Shen, B. and Dooner, H.K. (2001) The highly recombinogenic bz locus lies in an unusually gene-rich region of the maize genome. *Proc. Natl Acad. Sci. USA*, **98**, 8903–8908.
- Fu, H., Zheng, Z. and Dooner, H.K. (2002) Recombination rates between adjacent genic and retrotransposon regions in maize vary by two orders of magnitude. *Proc. Natl Acad. Sci. USA*, **99**, 1082–1087.
- García, V., Bruchet, H., Camescasse, D., Granier, F., Bouchez, D. and Tissier, A. (2003) AtATM is essential for meiosis and the somatic response to DNA damage in plants. *Plant Cell*, **15**, 119–132.
- García, V., Gray, S., Allison, R.M., Cooper, T.J. and Neale, M.J. (2015) Tel1 (ATM)-mediated interference suppresses clustered meiotic double-strand-break formation. *Nature*, **520**, 114–118.
- García, V., Phelps, S.E.L., Gray, S. and Neale, M.J. (2011) Bidirectional resection of DNA double-strand breaks by Mre11 and Exo1. *Nature*, **479**, 241–244.
- Girard, C., Crismani, W., Froger, N., Mazel, J., Lemhemdi, A., Horlow, C. and Mercier, R. (2014) FANCM-associated proteins MHF1 and MHF2, but not the other Fanconi anemia factors, limit meiotic crossovers. *Nucleic Acids Res.* **42**, 9087–9095.
- Giraut, L., Falque, M., Drouaud, J., Pereira, L., Martin, O.C. and Mézard, C. (2011) Genome-wide crossover distribution in *Arabidopsis thaliana* meiosis reveals sex-specific patterns along chromosomes. *PLoS Genet.* **7**, e1002354.
- Gore, M.A., Chia, J.-M., Elshire, R.J. et al. (2009) A first-generation haplotype map of maize. *Science*, **326**, 1115–1117.
- Grey, C., Barthès, P., Chauveau-Le Friec, G., Langa, F., Baudat, F. and de Massy, B. (2011) Mouse PRDM9 DNA-binding specificity determines sites of histone H3 lysine 4 trimethylation for initiation of meiotic recombination. *PLoS Biol.* **9**, e1001176.
- He, Y., Sidhu, G. and Pawlowski, W.P. (2013) Chromatin immunoprecipitation for studying chromosomal localization of meiotic proteins in maize. *Methods Mol. Biol.* **990**, 191–201.
- Hellsten, U., Wright, K.M., Jenkins, J., Shu, S., Yuan, Y., Wessler, S.R., Schmutz, J., Willis, J.H. and Rokhsar, D.S. (2013) Fine-scale variation in meiotic recombination in *Mimulus* inferred from population shotgun sequencing. *Proc. Natl Acad. Sci. USA*, **110**, 19478–19482.
- Higgins, J.D., Armstrong, S.J., Franklin, F.C.H. and Jones, G.H. (2004) The *Arabidopsis* MutS homolog AtMSH4 functions at an early step in recombination: evidence for two classes of recombination in *Arabidopsis*. *Genes Dev.* **18**, 2557–2570.
- Higgins, J.D., Buckling, E.F., Franklin, F.C.H. and Jones, G.H. (2008a) Expression and functional analysis of AtMUS81 in *Arabidopsis* meiosis reveals a role in the second pathway of crossing-over. *Plant J.* **54**, 152–162.
- Higgins, J.D., Vignard, J., Mercier, R., Pugh, A.G., Franklin, F.C.H. and Jones, G.H. (2008b) AtMSH5 partners AtMSH4 in the class I meiotic crossover pathway in *Arabidopsis thaliana*, but is not required for synapsis. *Plant J.* **55**, 28–39.

- Hinch, A.G., Tandon, A., Patterson, N. *et al.* (2011) The landscape of recombination in African Americans. *Nature*, **476**, 170–175.
- Hou, Y., Fan, W., Yan, L. *et al.* (2013) Genome analyses of single human oocytes. *Cell*, **155**, 1492–1506.
- Hudson, R.R. and Kaplan, N.L. (1985) Statistical properties of the number of recombination events in the history of a sample of DNA sequences. *Genetics*, **111**, 147–164.
- Hunt Morgan, T. (1916) *A Critique of the Theory of Evolution*. Princeton: Princeton University Press.
- Hyppa, R.W. and Smith, G.R. (2010) Crossover invariance determined by partner choice for meiotic DNA break repair. *Cell*, **142**, 243–255.
- Ines, O.Da., Degroote, F., Goubely, C., Amiard, S., Gallego, M.E. and White, C.I. (2013) Meiotic recombination in *Arabidopsis* is catalysed by DMC1, with RAD51 playing a supporting role. F. C. H. Franklin, ed. *PLoS Genet.* **9**, e1003787.
- Iyer, V. and Struhl, K. (1995) Poly(dA:dT), a ubiquitous promoter element that stimulates transcription via its intrinsic DNA structure. *EMBO J.* **14**, 2570–2579.
- Jeffreys, A.J., Kauppi, L. and Neumann, R. (2001) Intensely punctate meiotic recombination in the class II region of the major histocompatibility complex. *Nat. Genet.* **29**, 217–222.
- Joshi, N., Brown, M.S., Bishop, D.K. and Börner, G.V. (2015) Gradual implementation of the meiotic recombination program via checkpoint pathways controlled by global DSB Levels. *Mol. Cell*, **57**, 797–811.
- Kauppi, L., Jeffreys, A.J. and Keeney, S. (2004) Where the crossovers are: recombination distributions in mammals. *Nat. Rev. Genet.* **5**, 413–424.
- Kauppi, L., May, C.A. and Jeffreys, A.J. (2009) *Meiosis*. (Keeney, S. ed.). Totowa, NJ: Humana Press.
- Keeney, S., Giroux, C.N. and Kleckner, N. (1997) Meiosis-specific DNA double-strand breaks are catalyzed by Spo11, a member of a widely conserved protein family. *Cell*, **88**, 375–384.
- Khil, P.P. and Camerini-Otero, R.D. (2010) Genetic crossovers are predicted accurately by the computed human recombination map. *PLoS Genet.* **6**, e1000831.
- Khil, P.P., Smagulova, F., Brick, K.M., Camerini-Otero, R.D. and Petukhova, G.V. (2012) Sensitive mapping of recombination hotspots using sequencing-based detection of ssDNA. *Genome Res.* **22**, 957–965.
- Kleckner, N., Zickler, D., Jones, G.H., Dekker, J., Padmore, R., Henle, J. and Hutchinson, J. (2004) A mechanical basis for chromosome function. *Proc. Natl Acad. Sci. USA*, **101**, 12592–12597.
- Knoll, A., Higgins, J.D., Seeliger, K., Reha, S.J., Dangel, N.J., Bauknecht, M., Schröpfer, S., Franklin, F.C.H. and Puchta, H. (2012) The Fanconi anemia ortholog FANCM ensures ordered homologous recombination in both somatic and meiotic cells in *Arabidopsis*. *Plant Cell*, **24**, 1448–1464.
- Kurzbaue, M.-T., Uanschou, C., Chen, D. and Schlögelhofer, P. (2012) The recombinases DMC1 and RAD51 are functionally and spatially separated during meiosis in *Arabidopsis*. *Plant Cell*, **24**, 2058–2070.
- Lam, I. and Keeney, S. (2014) Mechanism and regulation of meiotic recombination initiation. *Cold Spring Harb. Perspect. Biol.* **7**, a016634.
- Lange, J., Pan, J., Cole, F., Thelen, M.P., Jasin, M. and Keeney, S. (2011) ATM controls meiotic double-strand-break formation. *Nature*, **479**, 237–240.
- Lenormand, T. and Dutheil, J. (2005) Recombination difference between sexes: a role for haploid selection. *PLoS Biol.* **3**, e63.
- Li, X., Li, L. and Yan, J. (2015) Dissecting meiotic recombination based on tetrad analysis by single-microspore sequencing in maize. *Nat. Commun.* **6**, 6648.
- Lichten, M. (2014) Tetrad, random spore, and molecular analysis of meiotic segregation and recombination. *Methods Mol. Biol.* **1205**, 13–28.
- Lichten, M. and Goldman, A.S. (1995) Meiotic recombination hotspots. *Annu. Rev. Genet.* **29**, 423–444.
- Liu, S., Yeh, C.-T., Ji, T., Ying, K., Wu, H., Tang, H.M., Fu, Y., Nettleton, D. and Schnable, P.S. (2009) Mu transposon insertion sites and meiotic recombination events co-localize with epigenetic marks for open chromatin across the maize genome. *PLoS Genet.* **5**, 13.
- Lu, S., Zong, C., Fan, W. *et al.* (2012) Probing meiotic recombination and aneuploidy of single sperm cells by whole-genome sequencing. *Science*, **338**, 1627–1630.
- Lynn, A., Soucek, R. and Börner, G.V. (2007) ZMM proteins during meiosis: crossover artists at work. *Chromosome Res.* **15**, 591–605.
- Macaisne, N., Novatchkova, M., Peirera, L., Vezon, D., Jolivet, S., Froger, N., Chelysheva, L., Grelon, M. and Mercier, R. (2008) SHOC1, an XPF endonuclease-related protein, is essential for the formation of class I meiotic crossovers. *Curr. Biol.* **18**, 1432–1437.
- Martini, E., Borde, V., Legendre, M., Audic, S., Regnault, B., Soubigou, G., Dujon, B. and Llorente, B. (2011) Genome-wide analysis of heteroduplex DNA in mismatch repair-deficient yeast cells reveals novel properties of meiotic recombination pathways. *PLoS Genet.* **7**, e1002305.
- de Massy, B. (2013) Initiation of meiotic recombination: how and where? Conservation and specificities among eukaryotes. *Annu. Rev. Genet.* **47**, 563–599.
- Mayer, K.F.X., Waugh, R., Brown, J.W.S. *et al.* (2012) A physical, genetic and functional sequence assembly of the barley genome. *Nature*, **491**, 711–716.
- McMahill, M.S., Sham, C.W. and Bishop, D.K. (2007) Synthesis-dependent strand annealing in meiosis. *PLoS Biol.* **5**, e299.
- McVean, G.A.T., Myers, S.R., Hunt, S., Deloukas, P., Bentley, D.R. and Donnelly, P. (2004) The fine-scale structure of recombination rate variation in the human genome. *Science*, **304**, 581–584.
- Melamed-Bessudo, C. and Levy, A.A. (2012) Deficiency in DNA methylation increases meiotic crossover rates in euchromatic but not in heterochromatic regions in *Arabidopsis*. *Proc. Natl Acad. Sci. USA*, **109**, E981–E988.
- Mercier, R., Jolivet, S., Vezon, D. *et al.* (2005) Two meiotic crossover classes cohabit in *Arabidopsis*: one is dependent on MER3, whereas the other one is not. *Curr. Biol.* **15**, 692–701.
- Mercier, R., Mézard, C., Jenczewski, E., Macaisne, N. and Grelon, M. (2014) The molecular biology of meiosis in plants. *Annu. Rev. Plant Biol.* **66**, 297–327.
- Mihola, O., Trachtulec, Z., Vlcek, C., Schimenti, J.C. and Forejt, J. (2009) A mouse speciation gene encodes a meiotic histone H3 methyltransferase. *Science*, **323**, 373–375.
- Mirouze, M., Lieberman-Lazarovich, M., Aversano, R., Bucher, E., Nicolet, J., Reinders, J. and Paszkowski, J. (2012) Loss of DNA methylation affects the recombination landscape in *Arabidopsis*. *Proc. Natl Acad. Sci. USA*, **109**, 5880–5885.
- Myers, S., Bottolo, L., Freeman, C., McVean, G. and Donnelly, P. (2005) A fine-scale map of recombination rates and hotspots across the human genome. *Science*, **310**, 321–324.
- Myers, S., Bowden, R., Tumian, A., Bontrop, R.E., Freeman, C., MacFie, T.S., McVean, G. and Donnelly, P. (2010) Drive against hotspot motifs in primates implicates the PRDM9 gene in meiotic recombination. *Science*, **327**, 876–879.
- Myers, S., Freeman, C., Auton, A., Donnelly, P. and McVean, G. (2008) A common sequence motif associated with recombination hot spots and genome instability in humans. *Nat. Genet.* **40**, 1124–1129.
- Neale, M.J., Pan, J. and Keeney, S. (2005) Endonucleolytic processing of covalent protein-linked DNA double-strand breaks. *Nature*, **436**, 1053–1057.
- Nicolas, A., Treco, D., Schultes, N.P. and Szostak, J.W. (1989) An initiation site for meiotic gene conversion in the yeast *Saccharomyces cerevisiae*. *Nature*, **338**, 35–39.
- Oliver, P.L., Goodstadt, L., Bayes, J.J. *et al.* (2009) Accelerated evolution of the Prdm9 speciation gene across diverse metazoan taxa. *PLoS Genet.* **5**, e1000753.
- Page, S.L. and Hawley, R.S. (2003) Chromosome choreography: the meiotic ballet. *Science*, **301**, 785–789.
- Pan, J., Sasaki, M., Kniewel, R. *et al.* (2011) A hierarchical combination of factors shapes the genome-wide topography of yeast meiotic recombination initiation. *Cell*, **144**, 719–731.
- Panizza, S., Mendoza, M.A., Berlinger, M., Huang, L., Nicolas, A., Shirahige, K. and Klein, F. (2011) Spo11-accessory proteins link double-strand break sites to the chromosome axis in early meiotic recombination. *Cell*, **146**, 372–383.
- Parvanov, E.D., Petkov, P.M. and Paigen, K. (2010) Prdm9 controls activation of mammalian recombination hotspots. *Science*, **327**, 835.
- Patterson, G.I., Kubo, K.M., Shroyer, T. and Chandler, V.L. (1995) Sequences required for paramutation of the maize B gene map to a region containing the promoter and upstream sequences. *Genetics*, **140**, 1389–1406.
- Pratto, F., Brick, K., Khil, P., Smagulova, F., Petukhova, G.V. and Camerini-Otero, R.D. (2014) Recombination initiation maps of individual human genomes. *Science*, **346**, 1256442.

- Qi, J., Chen, Y., Copenhaver, G.P. and Ma, H. (2014) Detection of genomic variations and DNA polymorphisms and impact on analysis of meiotic recombination and genetic mapping. *Proc. Natl Acad. Sci. USA*, **111**, 10007–10012.
- Qiao, H., Prasada Rao, H.B.D., Yang, Y. *et al.* (2014) Antagonistic roles of ubiquitin ligase HEI10 and SUMO ligase RNF212 regulate meiotic recombination. *Nat. Genet.* **46**, 194–199.
- Qin, J., Richardson, L.L., Jasin, M., Handel, M.A. and Arnheim, N. (2004) Mouse strains with an active H2-Ea meiotic recombination hot spot exhibit increased levels of H2-Ea-specific DNA breaks in testicular germ cells. *Mol. Cell. Biol.* **24**, 1655–1666.
- Qin, Y., Zhao, L., Skaggs, M.I. *et al.* (2014) ACTIN-RELATED PROTEIN6 regulates female meiosis by modulating meiotic gene expression in *Arabidopsis*. *Plant Cell*, **26**, 1612–1628.
- Reynolds, A., Qiao, H., Yang, Y. *et al.* (2013) RNF212 is a dosage-sensitive regulator of crossing-over during mammalian meiosis. *Nat. Genet.* **45**, 269–278.
- Rodgers-Melnick, E., Bradbury, P.J., Elshire, R.J., Glaubitz, J.C., Acharya, C.B., Mitchell, S.E., Li, C., Li, Y. and Buckler, E.S. (2015) Recombination in diverse maize is stable, predictable, and associated with genetic load. *Proc. Natl Acad. Sci. USA*, **112**, 3823–3828.
- Roitinger, E., Hofer, M., Köcher, T., Pichler, P., Novatchkova, M., Yang, J., Schögelhofer, P. and Mechtler, K. (2015) Quantitative phosphoproteomics of the ataxia telangiectasia-mutated (ATM) and ataxia telangiectasia-mutated and Rad3-related (ATR) dependent DNA damage response in *Arabidopsis thaliana*. *Mol. Cell Proteomics*, **14**, 556–571.
- Rowan, B.A., Patel, V., Weigel, D. and Schneeberger, K. (2015) Rapid and inexpensive whole-genome genotyping-by-sequencing for crossover localization and fine-scale genetic mapping. *G3: Genes - Genomes - Genetics*, **5**, 385–398.
- Saintenac, C., Falque, M., Martin, O.C., Paux, E., Feuillet, C. and Sourdille, P. (2009) Detailed recombination studies along chromosome 3B provide new insights on crossover distribution in wheat (*Triticum aestivum* L.). *Genetics*, **181**, 393–403.
- Salomé, P.A., Bomblies, K., Fitz, J., Laitinen, R.A.E., Warthmann, N., Yant, L. and Weigel, D. (2012) The recombination landscape in *Arabidopsis thaliana* F2 populations. *Heredity (Edinb)*, **108**, 447–455.
- Schwacha, A. and Kleckner, N. (1995) Identification of double Holliday junctions as intermediates in meiotic recombination. *Cell*, **83**, 783–791.
- Segal, E. and Widom, J. (2009) Poly(dA:dT) tracts: major determinants of nucleosome organization. *Curr. Opin. Struct. Biol.* **19**, 65–71.
- Séguéla-Arnaud, M., Crismani, W., Larchevêque, C. *et al.* (2015) Multiple mechanisms limit meiotic crossovers: TOP3 α and two BLM homologs antagonize crossovers in parallel to FANCM. *Proc. Natl Acad. Sci. USA* **112**, 4713–4718.
- Serrentino, M.-E., Chaplais, E., Sommermeyer, V. and Borde, V. (2013) Differential association of the conserved SUMO ligase Zip3 with meiotic double-strand break sites reveals regional variations in the outcome of meiotic recombination. *PLoS Genet.* **9**, e1003416.
- Shinohara, A., Ogawa, H. and Ogawa, T. (1992) Rad51 protein involved in repair and recombination in *S. cerevisiae* is a RecA-like protein. *Cell*, **69**, 457–470.
- Smagulova, F., Gregoret, I.V., Brick, K., Khil, P., Camerini-Otero, R.D. and Petukhova, G.V. (2011) Genome-wide analysis reveals novel molecular features of mouse recombination hotspots. *Nature*, **472**, 375–378.
- Sommermeyer, V., Béneut, C., Chaplais, E., Serrentino, M.E. and Borde, V. (2013) Spp1, a member of the Set1 complex, promotes meiotic DSB formation in promoters by tethering histone H3K4 methylation sites to chromosome axes. *Mol. Cell*, **49**, 43–54.
- Sun, H., Treco, D. and Szostak, J.W. (1991) Extensive 3'-overhanging, single-stranded DNA associated with the meiosis-specific double-strand breaks at the ARG4 recombination initiation site. *Cell*, **64**, 1155–1161.
- Sun, Y., Ambrose, J.H., Haughey, B.S. *et al.* (2012) Deep genome-wide measurement of meiotic gene conversion using tetrad analysis in *Arabidopsis thaliana*. *PLoS Genet.* **8**, e1002968.
- Szostak, J.W., Orr-Weaver, T.L., Rothstein, R.J. and Stahl, F.W. (1983) The double-strand-break repair model for recombination. *Cell*, **33**, 25–35.
- Thacker, D., Lam, I., Knop, M. and Keeney, S. (2011) Exploiting spore-autonomous fluorescent protein expression to quantify meiotic chromosome behaviors in *Saccharomyces cerevisiae*. *Genetics*, **189**, 423–439.
- The Tomato Genome Consortium (2012) The tomato genome sequence provides insights into fleshy fruit evolution. *Nature*, **485**, 635–641.
- Tiemann-Boege, I., Calabrese, P., Cochran, D.M., Sokol, R. and Arnheim, N. (2006) High-resolution recombination patterns in a region of human chromosome 21 measured by sperm typing. *PLoS Genet.* **2**, e70.
- Tischfield, S.E. and Keeney, S. (2012) Scale matters: the spatial correlation of yeast meiotic DNA breaks with histone H3 trimethylation is driven largely by independent colocalization at promoters. *Cell Cycle*, **11**, 1496–1503.
- Uanschou, C., Siwicz, T., Pedrosa-Harand, A. *et al.* (2007) A novel plant gene essential for meiosis is related to the human CtIP and the yeast COM1/SAE2 gene. *EMBO J.* **26**, 5061–5070.
- Venters, B.J. and Pugh, B.F. (2009) How eukaryotic genes are transcribed. *Crit. Rev. Biochem. Mol. Biol.* **44**, 117–141.
- Villeneuve, A.M. and Hillers, K.J. (2001) Whence meiosis? *Cell*, **106**, 647–650.
- Wei, F., Zhang, J., Zhou, S. *et al.* (2009) The physical and genetic framework of the maize B73 genome. *PLoS Genet.* **5**, 9.
- Wijeratne, A.J., Chen, C., Zhang, W., Timofejeva, L. and Ma, H. (2006) The *Arabidopsis thaliana* PARTING DANCERS gene encoding a novel protein is required for normal meiotic homologous recombination. *Mol. Biol. Cell*, **17**, 1331–1343.
- Wijner, E., Velikkakam James, G., Ding, J. *et al.* (2013) The genomic landscape of meiotic crossovers and gene conversions in *Arabidopsis thaliana*. *Elife*, **2**, e01426.
- Wu, T.C. and Lichten, M. (1994) Meiosis-induced double-strand break sites determined by yeast chromatin structure. *Science*, **263**, 515–518.
- Xu, X., Hsia, A.P., Zhang, L., Nikolau, B.J. and Schnable, P.S. (1995) Meiotic recombination break points resolve at high rates at the 5' end of a maize coding sequence. *Plant Cell*, **7**, 2151–2161.
- Yao, H. and Schnable, P.S. (2005) Cis-effects on meiotic recombination across distinct a1-sh2 intervals in a common *Zea* genetic background. *Genetics*, **170**, 1929–1944.
- Yelina, N.E., Choi, K., Chelysheva, L. *et al.* (2012) Epigenetic remodeling of meiotic crossover frequency in *Arabidopsis thaliana* DNA methyltransferase mutants. *PLoS Genet.* **8**, e1002844.
- Yelina, N.E., Ziolkowski, P.A., Miller, N., Zhao, X., Kelly, K.A., Muñoz, D.F., Mann, D.J., Copenhaver, G.P. and Henderson, I.R. (2013) High-throughput analysis of meiotic crossover frequency and interference via flow cytometry of fluorescent pollen in *Arabidopsis thaliana*. *Nat. Protoc.* **8**, 2119–2134.
- Young, J.A., Schreckhise, R.W., Steiner, W.W. and Smith, G.R. (2002) Meiotic recombination remote from prominent DNA break sites in *S. pombe*. *Mol. Cell*, **9**, 253–263.
- Zhang, X., Bernatavichute, Y.V., Cokus, S., Pellegrini, M. and Jacobsen, S.E. (2009) Genome-wide analysis of mono-, di- and trimethylation of histone H3 lysine 4 in *Arabidopsis thaliana*. *Genome Biol.* **10**, R62.