




AENSI Journals

Advances in Environmental Biology

Journal home page: <http://www.aensiweb.com/aeb.html>

Genetic Characterization of Shiga Toxin-producing *Escherichia coli* Strains Isolated from Frozen Bovine Meat in Algeria

¹Barka Mohammed Salih, ¹Benyoub Nour-Eddine, ²Henni Jamal-Eddine, ³Boudilmi Benabdallah and ²Kihal Mebrouk

¹Faculté des sciences de la nature de la vie de la terre et de l'univers, université d'Abou Bekr Belkaid Tlemcen Algérie.

²Laboratoire de microbiologie appliquée, département de biologie, faculté des sciences de la nature et de la vie, université d'Oran. Oran Es-Senia Algérie.

³Laboratoire Vétérinaire Régional Tlemcen Algérie.

ARTICLE INFO

Article history:

Received 22 October 2013

Received in revised form 14

January 2014

Accepted 20 January 2014

Available online 1 March 2014

Key words:

Escherichia coli O157:H7,

stx1, stx2, eae c, ehxA,

PCR, frozen bovine meat, Algeria.

ABSTRACT

In this study the presence of *E. coli* O157:H7 was investigated in 756 samples of frozen bovine meat imported from different countries using the standard method of culture and serotyping technique. The pathogen was detected in five (0.66 %) samples. All isolates were further characterized by PCR and serotyping. PCR showed that 1 isolate carried stx1 genes, 5 possessed stx2 genes and 1 both stx1 and stx2. Enterohemolysin (ehxA) and intimin (eae c) virulence genes were detected in 5 isolates and in 4 of the isolates respectively. The interaction with lactic acid bacteria revealed that all *E. coli* O157:H7 isolates were inhibited. Then all the isolates were tested using the method of KIRBY and BAUER according to the nccls standards to view resistance to antibiotics and was considered sensitive to the sixteen antimicrobial agents tested.

© 2014 AENSI Publisher All rights reserved.

To Cite This Article: Barka Mohammed Salih, Benyoub Nour-Eddine, Henni Jamal-Eddine, Boudilmi Benabdallah and Kihal Mebrouk., Genetic Characterization of Shiga Toxin-producing *Escherichia coli* Strains Isolated from Frozen Bovine Meat in Algeria. *Adv. Environ. Biol.*, 8(1), 6-13, 2014

INTRODUCTION

Escherichia coli enterohaemorrhagic (EHEC; also variously referred to as Verocytotoxin producing *E. coli*, VTEC and Shiga toxin producing *E. coli*, STEC) are an important group of food-borne pathogens that cause infections to humans in many parts of the world [35,3]. *E. coli* O157:H7 is the most important serotype causes simple diarrhea to the more complicated hemorrhagic colitis (HC) and hemolytic uremic syndrome (HUS) [14]. Cattle, especially the young ones, have been implicated as a principal source for *E. coli* O157:H7 [4]. Most infections caused by *E. coli* O157: H7 result from the consumption of contaminated and undercooked ground beef and unpasteurized fruit juices. Dairy products, cheese from raw milk [1,25].

Virulence factors include hemolysins and Shiga toxins (Stx) [13,33]. Members of family include *Stx1* and *Stx2* toxins encoded by *stx1* and *stx2* genes, respectively. Shiga toxins are the major virulence factors of EHEC which have cytotoxicity effects for human and animal eukaryotic cells and cause dangerous details like HUS and TTP [44] other virulence factor *eaec* and *ehxA* have been described [41,25]. In Algeria, no data were available until now on the fate and characteristics of STEC isolated from frozen bovine meat. The objective of the present study was to isolate *E. coli* O157:H7 from frozen bovine meat samples by conventional culture method and to confirm it by a serogroup-specific and PCR assay.

MATERIALS AND METHODS

Sample and bacterial isolates:

Seven hundred and fifty six samples of imported frozen bovine meat were analyzed. After received by the laboratory, enrichment cultures for each sample were carried out by combining 25 g of each sample with 225 ml of buffered peptone water supplemented into a stomacher bag, homogenized for at least 2 min and incubated at 37°C for 16–18 h. After incubation, the isolates belonging to *E. coli* were cultured on Sorbitol Mac Conkey medium added with MUG supplement (Biokar diagnostics) for detecting non sorbitol and MUG negative variants. This test required the identification guides to the pathogenic EHEC strains. The strains metabolizing

Corresponding Author: Barka Mohammed Salih, Faculté des sciences de la nature de la vie de la terre et de l'univers, université d'Abou Bekr Belkaid Tlemcen Algérie.

sorbitol are excluded, however the rest of the strains are tested for their immunological confirmation by the method of Farmer and Davis. Three *Lactobacillus* strains and one strain of *Leuconostoc* were used in this study. *Lactobacillus* strains were obtained from laboratory of applied microbiology, biology department, Oran university [6,36].

Biochemical and serological tests:

All the identified nonsorbitol fermenting colonies from the SMAC Agar were biochemically tested for confirmation by the API 20E, and subjected to slide agglutination with the the *E. coli* Latex kit O157:H7 (« Statens Serum Institute », Copenhagen, Denmark) and the agglutinating colonies were further processed for definite confirmation [23].

Interaction with lactic acid bacteria:

The research of the effect of bacteriocins produced by lactic acid bacteria is carried out by the method of the double layer [7]. Strains of *E. coli* O157: H7 inoculated on BHIB medium liquid are incubated 18 hours at 37 °C; while the lactic acid bacteria are inoculated in MRS spot on solid buffered in order to eliminate the effect of lactic acid (pH 5.4) then they were incubated 24 h at 30°C. After incubation of the *E. coli* O157: H7 were inoculated into BHIB semi solid Petri dishes are flooded with MRS strains to study and let incubated for 24 h at 37°C.

Antibiotic susceptibility test:

The isolated strains were tested for susceptibility to 16 antibiotics using the standard Kirby–Bauer method [8] and the following antibiotics discs were used: ampicillin (10 µg), chloramphenicol (30 µg), gentamycin (10 µg), colistin 10 µg, flumequin (30 µg), enrofloxacin (5 µg), nitrofurantoin (300 µg), cephalotin (30 µg), sulfamethoxazole-trimethoprim (1.25/23.75 µg), nalidixic acid (30 µg), norfloxacin (10 µg), ofloxacin (5 µg), kanamycin (30 µg), neomycin (30 µg), ceftiofur (30µg) and tetracycline (30 µg). Isolates were classed as sensitive or resistant to each antibiotic according to the Clinical and Laboratories Standards Institute (formally NCCLS) guidelines. *E. coli* strain ATCC 25922 and *S. aureus* strain ATCC 25923 were used for quality control.

Extraction of DNA:

The extraction of DNA was realized by the technique of phenol-chloroform. The bacteria are spread initially on MacConkey sorbitol, and then a clone is grown in 5 ml of LB medium. 1 ml of each culture was centrifuged 2500 g x 10 min, then the pellet was recovered and treated with lysis buffer (20% SDS, 10 mM Tris pH 7.5, 1M EDTA) and proteinase K at 10mg/ml. After heating at 56 °C for 2 hours, add the phenol-chloroform-isopropanol and follow the protocol of Sambrook *et al.* [45]. The DNA is recovered in 50 µl of sterile water.

PCR amplification:

The virulence genes of *E. coli* O157: H7 isolates were identified by polymerase chain reaction. *E. coli* ECL6611 were used as the positive control, *E. coli* ECL3463 and *E. coli* 29 as the negative control. The PCR reaction was performed in a 50 µl amplification mixture consisting of 5 µl 10 PCR buffer (500 mM KCl, 200 mM Tris-HCl), 8 µl dNTPs (10 mM), 1.5 µl MgCl₂ (50 mM), 2.5 µl of each primer (10 µM), 0.5 µl of Taq DNA polymerase (5 Unit/µl) and 1 µl of template [40,43]. The primers used in this PCR are presented in the following table:

Table 1: Oligonucleotide primers sequences used in PCR (*stx1a* and *stx2b* for shigatoxine, *ehxA* for enterohemolysin and *eae c* for intimin).

Gene	Primer	Oligonucléotide	séquence (5'–3')	Fragment size (bp)	Tm	Référence
<i>stx1a</i>	VT1-A	CGCTGAATGTCATTCGCTCTGC	302	71.2		Blanco <i>et al.</i> (2003)
	VT1-B	CGTGGTATAGCTACTGTCACC	58.5			
<i>stx2b</i>	VT2-A	CTTCGGTATCCTATTCCCGG	516	64.2		Blanco <i>et al.</i> (2003)
	VT2-B	CTGCTGTGACAGTGACAAAAACGC	69.1			
<i>ehxA</i>	HlyA1	GGTGCAGCAGAAAAAGTTGTAG	1551	62.7		Schmidt <i>et al.</i> (1995)
	HlyA4	TCTCGCCTGATAGTGTGGTA	63.4			
<i>eae c</i>	EAE-1	GAGAAATGAAATAGAAGTCGT	775	52.4		Blanco <i>et al.</i> (2003)
	EAE-2	GCGGTATCTTTCGCGTAATCGCC	72.6			

The amplifications were performed for:

stx1 at 95°C 30 s, 30 cycles of 95°C for 15 s; 52°C for 60s; 72°C for 90s.

stx2 at 95°C 30 s, 30 cycles of 95°C for 15 s; 55°C for 30 s; 72°C for 90s.

eae c at 95°C 30 s, 30 cycles of 95°C for 15 s; 45°C for 60 s; 72°C for 90s.

ehxA at 95°C 30 s, 30 cycles of 95°C for 15 s; 55°C for 60 s, 72°C for 90s.

Both positive and negative control reactions were included in each PCR amplification experiment.

Electrophoresis of PCR products:

PCR products were separated by electrophoresis on 1, 5 % (w/v) agarose gel (Invitrogen) at 100 V for 40 min in Tris acetate buffer. The gels were stained in ethidium bromide, illuminated by UV-transilluminator and documented by a gel documentation apparatus. Each gel contained a 1 kb DNA molecular weight marker (Fermentas, USA).

*Results:**Isolation and identification of E. coli O157:H7:*

From 756 samples of meat analyzed; 3 *Escherichia coli* sorbitol negative were isolated and identified from Uruguay samples, 1 from Brazil and 1 from Argentina (Tab. 2)

Then the strains are identified by the API 20E system that enables us to confirm the sorbitol negative test and especially to have the chemotype of these strains.

One strain has chemotype 5144112, in fact we find that this strain in addition to sorbitol, it does not ferment sucrose, melibiose and amygdalin. While the four other strains have the same chemotype 5144172.

Table 2: Number of *E. coli* O157:H7 detected in imported frozen meat in Algeria.

Country	Uruguay	Brazil	Argentina	Ireland	Australia	New Zealand	Total
Number of Samples examined	365	243	105	21	16	06	756
Number of positive samples (%)	3(0.39)	1(0.13)	1(0.13)	0(0.0)	0(0.0)	0(0.0)	5

The identification of the strains was followed by the procedure applied by Barka and Kihal [6].

Interaction with lactic acid bacteria:

All strains were inhibited in contact with lactic acid bacteria used. Figure (1) shows the zones of inhibition




Fig. 1: Inhibition effect of lactic acid bacteria A4 : *Lactobacillus johnsonii*, 5 : *Lactobacillus plantarum*, 58: *Lactobacillus paracasei*, 68: *Lactobacillus plantarum*, leu : *Leuconostoc*, toward *E. coli* strains 26 and 28.

Antibiotic susceptibility test:

Antibiotic susceptibility profiles indicated that only one resistance strain to tetracycline was observed, the strain number 25.

Genetic methods for detecting E. coli O157: H7:

In this study all strains possess the genes *stx2* and *ehxA*, one strain (number 27) does not possess the *eae* gene but instead it is the only strain that has the gene *stx1*. Table 3 summarizes the results of pcr and Figures (2, 3, 4 and 5) show the presence of bands corresponding to each gene on the electrophoresis gels

Table 3: Genes detected in the different strains *E. coli* O157: H7 isolated from frozen bovin meat in Algeria.

Gène Strain of <i>E. coli</i>	<i>Stx1</i>	<i>Stx2</i>	<i>Stx1 et Stx2</i>	<i>eae</i>	<i>ehxA</i>
25	-	+	-	+	+
26	-	+	-	+	+
27	+	+	+	-	+
28	-	+	-	+	+
30	-	+	-	+	+
<i>E. coli</i> ECL6611	+	+	+	+	+
<i>E. coli</i> ECL3463	-	-	-	-	-




Fig. 2: Amplification products of *stx1* gene by polymerase chain reaction (PCR). M (1 kb), *E. coli* ECL6611 strain is the positive control. The negative control is the strain of *E. coli* ECL3463. Only strain 27 has the *stx1* gene.




Fig. 3: Amplification products of *stx2* gene by polymerase chain reaction (PCR). M (1 kb), *E. coli* ECL6611 strain is the positive control. The negative control is the strain of *E. coli* ECL3463.. The strains 25, 26, 27, 28 and 30 have the *stx2* gene.




Fig. 4: Amplification products of *eae* gene by polymerase chain reaction (PCR). M (1 kb), *E. coli* ECL6611 strain is the positive control. The negative control is the strain of *E. coli* ECL3463. The strains 25, 26, 28 and 30 have the *eae* gene.




Fig. 5: Amplification products of *ehxA* gene by polymerase chain reaction (PCR). M (1kb), *E. coli* ECL6611 strain is the positive control. The negative control is the strain of *E. coli* ECL3463. The strains 25, 26, 27, 28 and 30 have the *ehxA* gene.

Discussion:

Approximately 93% of strains of *E. coli* of human origin were sorbitol positive in 24 hours and are β -glucuronidase positive, in contrast, *E. coli* O157: H7 does not ferment sorbitol and are β -glucuronidase negative.

The absence of sorbitol fermentation, for example, justified the use of sorbitol MacConkey agar (SMAC), which has undergone several changes in the objective of increasing the selectivity against O157: H7. The biochemical methods are interesting and fundamental for the detection of *E. coli* O157: H7. Beutin *et al.* [9].

From 756 samples tested, five *E. coli* O157: H7 strains were isolated with a rate of 0.66%, the rate of *E. coli* O157: H7 varies from one country to another. The only known study on *E. coli* O157: H7 in Algeria was made in 2006 by Chahed after analyzing 230 samples of beef carcasses from slaughterhouse in the region of Algiers in which had isolated two *E. coli* O157: H7. In France Vernozy-Rozand *et al.*, [53] have detected a rate of 0.12% in

the meat industry and 1.9% of carcasses [42], in Switzerland no *E. coli* O157: H7 were detected from 400 samples [24]. 1% of the samples were positive for *E. coli* O157: H7 in Spain [5], 1% and 0.4% in England [17], 1% in the Netherlands [27] and 0.4% in Italy [21,50] and 0.3% in the Czech Republic [29].

Other studies have revealed much higher rates including one made in Argentina with 3.8% [20], Peru 19% [32] and Jordan with 7.8% in carcasses [39].

According to Nastasijevic *et al.*, [34] the presence of *E. coli* O157: H7 is usually the result of contamination during the slaughter process. It was previously reported that this contamination is correlated with the presence of STEC in feces and hides [22].

Concerning the interaction with lactic bacteria some studies have shown inhibition of gram-negative bacteria by bacteriocins [51].

Currently to prevent food contamination by pathogenic bacteria in the case of *E. coli* O157: H7, some authors recommend to use in addition to the barriers used in food safety which are temperature, water activity (Aw), pH, redox potential (Eh), vacuum packaging, modified atmosphere, high hydrostatic pressure (HHP), UV and competitive flora (LAB producing antimicrobial compounds), the combined use of bacteriocins produced by the plant competitive with storage methods listed above to create a series of obstacles during the manufacturing process to reduce food spoilage. Indeed, these different barriers positively influence the activity of many bacteriocins by increasing the permeability of cell membranes of target bacteria [19,2].

Antibiotic susceptibility shows result different of other studies around the world or a fairly high rate of resistance was observed. Recent studies have revealed an increasing trend of *E. coli* O157: H7 against antibiotics [31,49].

For example, in 2005, about 35% of *E. coli* O157: H7 isolated from meat and meat products in Gabone, Botswana, were resistant to cephalothin, sulfatriad, colistin sulfate and tetracycline [31]. In Korea one study found a rate of 71% resistance to at least one antibiotic and is the highest rate was reported today [28]. An American study has shown a resistance rate of 20% to tetracycline and 14% to sulfonamides [48].

In our study, we found that *stx2* was the predominant gene over *stx1* and all the strains possess the *ehxA*. It can be concluded that all strains carry one or more genes in pathogenicity. These results are similar to those obtained by Blanco *et al.* [11,12], 0% of cows, calves 0.6%, but less than obtained by Oporto *et al.*, [38] in calves 6.7% and Sánchez *et al.*, [46] 7 strains of 268 samples analyzed.

Khan in 2002 in India found 44.5% of strains of *E. coli* O157: H7 bearing both the *stx1* and *stx2*. The genes *stx2* and *eaeA* are more frequently isolated than *stx1* in most studies in the United States, Japan and European countries [30;15]. In a U.S.A study the gene *stx1* have been detected in any sample, while 22% of isolates carried the *stx2* gene [39].

Some studies show that the presence of the *stx2* gene was most of the time associated with the presence of the gene *eaeA* [37]. Only some studies have shown the presence in the same strain genes *stx1*, *stx2* and *eaeA* [26,52].

In our study the genes *stx2* and *eaeA* were detected in most strains analyzed. This is important because the presence of these genes is strongly associated with human disease [46].

These results show that there may be contamination of frozen meat must done research of these pathogen bacteria before the placing on the market.

REFERENCES

- [1] Allerberger, F., M. Wagner, P. Schweiger, H.P. Rammer, A. Resch, M.P. Dierich, A.W. Friedrich and H. Karch, 2001. *Escherichia coli* O157 infections and unpasteurised milk. *Euro. Surveill.*, 6: 147-151.
- [2] Ananou, S., A. Muñoz, M. Martínez-Bueno, P. González-Tello, A. Gálvez, M. Maqueda and E. Valdivia. 2009. Evaluation of an enterocin AS-48 enriched bioactive powder obtained by spray drying. *Food Microbiol.*, 27(1): 58-63.
- [3] Armstrong, G.L., J. Hollingsworth and J.R. Morris, 1996. Emerging foodborne pathogens: *Escherichia coli* O157:H7 as a model of entry of a new pathogen into the food supply of the developed world. *Epidemiol. Rev.*, 18: 29-51.
- [4] Ateba, C.N. and M. Mbeve, 2013. Determination of the genetic similarities of fingerprints from *Escherichia coli* O157:H7 isolated from different sources in the North West Province, South Africa using ISR, BOXAIR and REP-PCR analysis. *Microbiol Res.*, 168(7): 438-46.
- [5] Azucena, M., B. Miguel, E.B. Jesús, D. Ghizlane, L. Cecilia, J. Paula, P.A. María, E. Aurora, I.B. María, A.G. Enrique and B. Jorge, 2007. Serotypes, virulence genes and intimin types of Shiga toxin (verocytotoxin)-producing *Escherichia coli* isolates from minced beef in Lugo (Spain) from 1995 through 2003. *BioMed. Cent. Microbiol.*, 7: 13.
- [6] Barka, M.S. M. Kihal, 2010. Prevalence of *Escherichia coli* enterohemorrhagic O157:h7 in Frozen Bovine Meat in Algeria. *J. Appl. Sci. Res.*, 6(11): 1576-1580.
- [7] Barefoot, S.F. and T.R. Klaenhammer, 1983. Detection and activity of lacticin B, has bacteriocin produced by *Lactobacillus acidophilus*. *Appl. Environ. Microbiol.*, 45(6): 1808-1815.

- [8] Bauer, A.W., W. Kirby, J.C. Sherris and M. Truck, 1966. Antibiotic susceptibility testing by a standardized single disc method. *Am. J. Clin. Pathol.*, 45: 493-496.
- [9] Beutin, L., A. Miko, G. Krause, K. Pries, S. Haby, K. Steege and N. Albrecht, 2007. Identification of human-pathogenic strains of Shiga toxin-producing *Escherichia coli* from food by a combination of serotyping and molecular typing of Shiga toxin genes. *Appl. Environ. Microbiol.*, 73: 4769-4775.
- [10] Bonardi, S., E. Maggi, A. Bottarelli, M.L. Pacciarini, A. Ansuini, G. Vellini, S. Morabito and A. Caprioli, 1999. Isolation of verocytotoxin-producing *Escherichia coli* O157:H7 from cattle at slaughter in Italy. *Vet. Microbiol.*, 67: 203-211.
- [11] Blanco, M., J.E. Blanco, J. Blanco, E.A. González, A. Mora, C. Prado, L. Fernández, M. Rio, J. Ramos and M.P. Alonso, 1996. Prevalence and characteristics of *Escherichia coli* serotype O157:H7 and other verotoxin-producing *E. coli* in healthy cattle. *Epidemiol. Inf.*, 117: 251-257.
- [12] Blanco, M., J.E. Blanco, J. Blanco, A. Mora, C. Prado, M.P. Alonso, M. Mouriño, C. Madrid, C. Balsalobre and A. Juárez, 1997. Distribution and characterization of faecal verotoxin-producing *Escherichia coli* (VTEC) isolated from healthy cattle. *Vet. Microbiol.*, 54: 309-319.
- [13] Brooks, G.F., J.S. Butel and S.A. Morse, 1998. *Jawetz, Melnick & Adelberg's Medical Microbiology*, pp: 218-230. Appleton & Lange.
- [14] Browning, N.G., J.R. Botha and J. Sacho, 1990. *Escherichia coli* O157:H7 haemorrhagic colitis. Report of the first South African case. *South African Journal of Surgery*, 28: 28-29.
- [15] Brusa, V., V. Aliverti, F. Aliverti, E.E. Ortega, J.H. de la Torre, L.H. Linares, M.E. Sanz, A.I. Etcheverría, N.L. Padola, L. Galli, P. Peral García, J. Copes and G.A. Leotta, 2012. Shiga toxin-producing *Escherichia coli* in beef retail markets from Argentina. *Front Cell Infect Microbiol.*, 2: 171.
- [16] Chahed, A., B. China, J. Mainil and G. Daube, 2006. Prevalence of enterohaemorrhagic *Escherichia coli* from serotype O157 and other attaching and effacing *Escherichia coli* on bovine carcasses in Algeria. *J. Appl. Microbiol.*, 101(2): 361-8.
- [17] Chapman, P.A., D.J. Wright, R. Higgins, 1993. Untreated milk as a source of verotoxigenic *Escherichia coli* O157. *Vet. Rec.*, 133: 171-172.
- [18] Chapman, P.A., C.A. Siddons, A.T. Cerdan Malo and M.A. Harkin, 2000. A one year study of *Escherichia coli* O157 in raw beef and lamb products. *Epidemiol. Infect.*, 124: 207-213.
- [19] Chawla, S.P. and R. Chander, 2004. Microbiological safety of shelf-stable meat products prepared by employing hurdle technology. *Food Control.*, 15: 559-563.
- [20] Chinen, I., J.D. Tanaro, E. Miliwebsky, L.H. Lound, G. Chillemi, S. Ledri, A. Baschkier, M. Scarpin, E. Manfredi and M. Rivas, 2001. Isolation and characterization of *Escherichia coli* O157:H7 from retail meats in Argentina. *J. Food Prot.*, 64: 1346-1351.
- [21] Conedera, G., P. Dalvit, M. Martín, G. Galiero, M. Gramaglia, E. Goffredo, G. Loffredo, S. Morabito, D. Ottaviani, F. Paterlini, G. Pezzotti, M. Pisanu, P. Semprini and A. Caprioli, 2004. Verocytotoxin-producing *Escherichia coli* O157 in minced beef and dairy products in Italy. *Int. J. Food Microbiol.*, 96: 67-73.
- [22] Elder, R.O., J.E. Keen, G.R. Siragusa, G.A. Barkocy-Gallagher, M. Koohmaraie and W.W. Laegreid, 2000. Correlation of enterohemorrhagic *Escherichia coli* O157 prevalence in feces, hides, and carcasses of beef cattle during processing. Proceedings of the National Academy of Sciences of the United States of America, 97: 2999-3003.
- [23] Farmer, J. and B. Davis, 1985. H7 antiserum sorbitol fermentation medium: a single tube screening medium for detecting *Escherichia coli* O157:H7 associated with hemorrhagic colitis. *J. Clin. Microbiol.*, 22: 620-625.
- [24] Fantelli, K. and R. Stephan, 2001. Prevalence and characteristics of Shigatoxin-producing *Escherichia coli* and *Listeria monocytogenes* strains isolated from minced meat in Switzerland. *Int. J. Food Microbiol.*, 70: 63-69.
- [25] Ghanbarpour, R. and E. Oswald, 2010. Phylogenetic distribution of virulence genes in *Escherichia coli* isolated from bovine mastitis in Iran. *Res. Veter. Sci.*, 88: 6-10.
- [26] Guyon, R., F. Dorey, J.P. Malas, F. Grimont, J. Foret, B. Rouviere and J.F. Collobert, 2001. Superficial contamination of bovine carcasses by *Escherichia coli* O157:H7 in a slaughterhouse in Normandy (France). *Meat Sci.*, 58: 329-331.
- [27] Heuvelink, A.E., Zwartkruis-Nahuis, J.T., Beumer, R.R. and E. de Boer, 1999. Occurrence and survival of verocytotoxin-producing *Escherichia coli* O157 in meats obtained from retail outlets in The Netherlands. *J. Food Prot.*, 62: 1115-1122.
- [28] Ju-Yeon, Y., M. Bo-Mi, O.H. In-Gyeong, K. Byeong, B. Li-Guang, L. Byeong-Su, K. Barry, D. Stein and H.L. John, 2006. Antimicrobial resistance of *Escherichia coli* O157 from cattle in Korea. *Int. J. Food Microbiol.*, 106: 74-78.
- [29] Lukášová, J., B. Abraham and S. Cupáková, 2004. Occurrence of *Escherichia coli* O157 in raw material and food in Czech Republic. *J. Vet. Med. B.*, 51: 77-81.

- [30] Lim, J.Y., J.W. Yoon and C.J. Hovde, 2010. A brief overview of *Escherichia coli* O157:H7 and its plasmid O157, *J. Microbiol. Biotechnol.*, 20: 1-10.
- [31] Magwira, C.A., B.A. Gashe and E.K. Collison, 2005. Prevalence and antibiotic resistance profiles of *Escherichia coli* O157:H7 in beef products from retail outlets in Gaborone, Botswana. *J. Food Prot.*, 68(2): 403-406.
- [32] Mora, A., S.L. León, M. Blanco, J.E. Blanco, C. López, G. Dahbi, A. Echeita, E.A. González and J. Blanco, 2007. Phage types, virulence genes and PFGE profiles of Shiga toxin-producing *Escherichia coli* O157:H7 isolated from raw beef meat, soft cheese and vegetables in Lima (Peru). *Int. J. Food Microbiol.*, 114: 204-210.
- [33] Murray, P.R., E.J. Baron, M.A. Pfaller, F.C. Tenover and R.H. Tenover, 1999. Manual of Clinical Microbiology. *American Society for Microbiology*, pp: 450-452.
- [34] Nastasišević, I., R. Mitrović and S. Buncić, 2009. The occurrence of *Escherichia coli* O157 in/on faeces, carcasses and fresh meats from cattle. *Meat Science*, 82: 101-105.
- [35] Nataro, J.P. and J.B. Kaper, 1998. Diarrheagenic *Escherichia coli*. *Clin. Microbiol. Rev.*, 11: 142-201.
- [36] Ngwa, G.A., R. Schop, S. Weir, C.G. León-Velarde and J.A. Odumeru, 2013. Detection and enumeration of *E. coli* O157:H7 in water samples by culture and molecular methods. *J. Microbiol. Met.*, 92: 164-172.
- [37] Omisakin, F., M. Macrae, I.D. Ogden and N.J. Strachan, 2003. Concentration and prevalence of *Escherichia coli* O157 in cattle feces at slaughter. *Appl. Environ. Microbiol.*, 69: 2444-2447.
- [38] Oporto, B., J.I. Esteban, G. Aduriz, R.A. Juste and A. Hurtado, 2008. *Escherichia coli* O157:H7 and non-O157 Shiga toxin-producing *E. coli* in healthy cattle, sheep and swine herds in Northern Spain. *Zoonoses and Public Health*, 55(2): 73-81.
- [39] Osaili, T.M., A.R. Alaboudi and M. Rahalah, 2012. Prevalence and antimicrobial susceptibility of *Escherichia coli* O157:H7 on beef cattle slaughtered in Amman abattoir. *Meat Science*, 93: 463-468.
- [40] Padola, N.L., M.E. Sanz, J.E. Blanco, M. Blanco, J. Blanco, A.I. Etcheverría, G.H. Arroyo, M.A. Usera and A.E. Parma, 2004. Serotypes and virulence genes of bovine Shiga toxin-producing *Escherichia coli* (STEC) isolated from a feedlot in Argentina. *Vet. Microbiol.*, 100: 3-9.
- [41] Pilpott, D. and F. Ebel, 2003. *E. coli: shiga toxin methods and protocols*. 1st. Edn., Totowa, Humana Press Inc., pp: 9-45.
- [42] Rodgers, S., 2001. Preserving non-fermented refrigerated foods with microbial cultures: a review. *Trends Food Sci. Technol.*, 12: 276-284.
- [43] Rubén Gordillo, J., J. Córdoba, J. María, M. Andrade, I. Luque and M. Rodríguez, 2011. Development of PCR assays for detection of *Escherichia coli* O157:H7 in meat products. *Meat Science*, 88: 767-773.
- [44] Saif, Y.M., H.J. Barnes, J.R. Glisson, A.M. Fadly, L.R. McDougald and D.E. Swayne, 2003. *Diseases of Poultry*, pp: 631-644. Iowa State University Press.
- [45] Sambrook, J., E.F. Fritsch and T. Maniatis, 1989. *Molecular Cloning : A Laboratory Manual* 2nd Edition, Vol. 3, pages E3 -E4; Cold Spring Harbor Laboratory Press, Washington, D.C., USA.
- [46] Sánchez, S., R. Martínez, A. García, J. Blanco, A. Echeita, J. Hermoso de Mendoza, J. Rey and J.M. Alonso, 2010a. Shiga toxin-producing *Escherichia coli* O157:H7 from extensive cattle of the fighting bulls breed. *R. in. Vet. Sci.*, 88: 208-210.
- [47] Sánchez, S., R. Martínez, A. García, D. Vidal, J. Blanco, M. Blanco, J.E. Blanco, A. Mora, S. Herrera-Leon, A. Echeita, J.M. Alonso and J. Rey, 2010b. Detection and characterisation of O157:H7 and non-O157 Shiga toxin-producing *Escherichia coli* in wild boars. *Vet. Microbiol.*, 143: 420-423.
- [48] Schroeder, C.M., C. Zhao, C. DeRoy, J. Torcolini, S. Zhao, D.G. White, D.D. Wagner, P.F. McDermott, R.D. Walker and J. Meng, 2002. Antimicrobial resistance of *Escherichia coli* O157 isolated from humans, cattle, swine, and food. *Appl. Environ. Microbiol.*, 68: 576- 581.
- [49] Solomakos, N., A. Govaris, A.S. Angelidis, S. Pournaras, A. Burriel, R. Kritas and S.K. Papageorgiou, 2009. Occurrence, virulence genes and antibiotic resistance of *Escherichia coli* O157 isolated from raw bovine, caprine and ovine milk in Greece. *Food Microbiol.*, 26: 865-871.
- [50] Stampi, S., A. Caprioli, G. De Luca, P. Quaglio, R. Sacchetti and F. Zanetti, 2004. Detection of *Escherichia coli* O157 in bovine meat products in northern Italy. *Int. J. Food Microbiol.*, 90: 257-262.
- [51] Todorov, S.D. and L.M. Dicks, 2004. Screening of lactic-acid bacteria from South African barley beer for the production of bacteriocin-like compounds. *Folia Microbiol. (Praha)*, 49: 406-10.
- [52] Tuteneš, A.V., D. Pierard, J. Van Hoof, M. Cornelis and L.D. Zutter, 2003. Isolation and molecular characterization of *E. coli* O157 isolated from cattle, pigs and chickens at slaughter. *Int. J. Food Microbiol.*, 84: 63-69.
- [53] Vernozy-Rozand, C., J. Bouvet, M.P. Montet, C. Bavai, S. Ray-Gueniot, C. Mazuy-Cruchaudet and Y. Richard, 2002. Survey of retail raw milk cheeses for Verotoxin-producing *E. coli* (VTEC) and *E. coli* O157:H7 in France. In 102th General Meeting of American Society for Microbiology May, 19-20 Salt-Lake City, USA.