



## Using music therapy protocols in the treatment of premature infants: An introduction to current practices

Lori F. Gooding, PHD, MT-BC (NICU-MT)\*

The Florida State University, College of Music, 122N Copeland Street, Tallahassee, FL 32306-1180, United States

### ARTICLE INFO

**Keywords:**  
Music therapy  
Infant  
Multimodal  
Medical  
Premature  
PAL

### ABSTRACT

Medical music therapy, which has developed rapidly in the last 15 years, has been effectively used with preterm infants in neonatal intensive care units. The purpose of this article is to provide an introduction to current music therapy protocols for premature infants, as well as highlight research supporting the use of these procedures to address a variety of medical and developmental objectives. Protocols covered include (a) music combined with kangaroo care, (b) multimodal stimulation, (c) developmentally appropriate music listening and (d) pacifier activated lullabies (PAL). Research suggests that these procedures have been effective in addressing a wide range of medical and developmental needs. Positive outcomes include: (a) reduced length of stay, (b) stabilized oxygen saturation levels, (c) increased stimulation tolerance, (d) reduced stress-related behaviors, (e) enhanced parent–infant bonding and (f) improved parent–child interactions.

© 2010 Elsevier Inc. All rights reserved.

### Introduction

Because of advances in medical care in the 1980s and 1990s, premature infants now have higher survival rates than infants born in previous decades (Wilson-Costello, Friedman, Minich, Fanaroff, & Hack, 2005). Today, infants as young as 23 weeks gestational age are surviving (17% survival rate), with survival rates increasingly dramatically for infants 24 gestational weeks and older. In fact, infants born at 24 weeks have a survival rate of 68%, infants born at 25 weeks have a survival rate of 71% and infants born at 26 weeks gestational age have an 80% survival rate (Gibson, Carney, & Wales, 2006; Hoekstra, Ferrara, Couser, Payne, & Connett, 2004). Furthermore, premature infants born between 35 and 38 weeks gestational age now have a 98% survival rate (Kelly, 2006).

Along with increased survival rates comes increased risk of significant neurodevelopmental impairment (Gibson et al., 2006; Wilson-Costello et al., 2005). Evidence suggests that significant growth impairments are almost completely universal in extremely premature infants (Gibson et al., 2006). Furthermore, research suggests that impairments are most significant for infants born at 23 weeks gestational age; these infants are more likely to be classified as “severely” impaired while infants born at 24–26 weeks gestational age are more likely to be classified as “normal” (Hoekstra et al., 2004). Additionally, follow-up research conducted with chil-

dren who were born prematurely suggests that higher rates of asthma, cerebral palsy, subnormal cognitive functioning, poor academic achievement and behavioral problems are common, and that impairments may lead to an increased predisposition to morbidity and premature mortality in adulthood (Gibson et al., 2006; Hack, 2007).

When examining specific neurodevelopmental impairments in preterm infants, a wide range of issues can be found. Infants born prior to 35 weeks do not have the ability to suck and swallow and rely on gastric lavage feeding to survive. Moreover, premature infants have difficulty processing nutrients, which leads to impaired growth development (Gibson et al., 2006). Research suggests that preterm infants have smaller brain volume and more limited cognitive abilities (Standley, 2003a, 2003b). Research also suggests that medical treatment for premature infants may cause long-term effects. For example, infants may develop visual impairments due to high levels of oxygen or permanent hearing loss may occur as a result of the use of ototoxic drugs (Standley, 2003a, 2003b). Additionally, premature infants may develop hyperactivity, learning disabilities or attention deficit disorders. It is believed that these neurobiological conditions may be the result of persistently high stress levels during medical treatments (Standley, 2003a, 2003b).

In addition to neurodevelopmental issues, preterm infants often have medical complications. Jaundice and intraventricular hemorrhage are all fairly common in preterm infants (Standley, 2003a, 2003b). Gastroesophageal reflux (GER) may also be an issue for preterm infants; in fact, as many as 50% of all infants between birth and 3 months deal with GER, with premature birth serving

\* Correspondence address: 466 Beechwood Dr., Evans, GA 30809, United States. Tel.: +1 850 241 2533.

E-mail address: cgood830@aol.com.

as one of the predisposing factors (Kelly, 2006). Preterm infants are also at risk for respiratory problems, including susceptibility to Respiratory Syncytial virus and Influenza (Kelly, 2006). Finally, preterm infants are at risk for developing apnea of prematurity, which can be defined as “sudden cessation of breathing that lasts for at least 20 s or is accompanied by bradycardia and oxygen desaturation in an infant less than 37 weeks’ gestational age” (Kelly, 2006, p. 298). Apnea of prematurity often occurs until the infant is capable of temperature regulation and able to fully nipple feed (Kelly, 2006).

A variety of treatments have been utilized with preterm infants in neonatal intensive care units (NICU) to address medical and developmental needs. One promising intervention now being utilized in NICUs is music therapy. According to Standley (2003a, 2003b), music therapy is a “research based profession with specific a priori objectives to facilitate medical, psychological or educational goals” (p. 64). Medical music therapy, which has developed rapidly in the last 15 years, has been shown to be effective in reducing stress, reducing the perception of pain, reducing stimulus deprivation and promoting psychological adjustment to trauma. Just as importantly, music has few, if any, unwanted side effects (Standley, 2003a, 2003b).

### Sound and hearing

In order to understand how music can be applied therapeutically with premature infants, one must first understand the basic principles of sound and hearing development in infants. Sound is vibratory energy (waves) transmitted through a medium (air, water, etc.). Sound waves move via compressions and rarefactions; the number of compressions and rarefactions per second is known as Hertz (Hz). Human beings can hear sounds that range from 20 to 20,000 Hz, with the greatest sensitivity found between 1000 and 4000 Hz (Lasky & Williams, 2005). However, Hepper and Shahidullah (1994) suggest that infant hearing is restricted to 500–1000 Hz, with considerable expansion occurring after birth.

Sound waves are transmitted through the hearing apparatus and converted from mechanical energy into electrochemical energy. Sound waves are transmitted through the fluid filled cochlea in the inner ear, with lower frequencies peaking near the apex and higher frequencies peaking near the base. Outer hair cells in the cochlea are bent through a shearing motion caused by displacement. This in turn bends the inner hair cells, opening potassium channels. The inner hair cells become polarized, sending transmissions along the auditory pathway to and from the brain via the VIIIth cranial nerve, which is completely myelinated by the 5th month post-conception (Lasky & Williams, 2005).

Auditory functioning begins after the 20th week of gestation and cochlea maturation occurs by approximately 35 weeks gestational age (Lasky & Williams, 2005). Even though auditory development is complete around 35 weeks gestational age, fetal response to sound occurs much earlier during development. Hepper and Shahidullah (1994) detected responses as early as 19 weeks post-menstruation age (PMA is a slightly different calculation than the gestational age based on date of mother’s last menses) to a 500 Hz tone. Others have found responses between 20 and 25 weeks PMA (Lasky & Williams, 2005). In fact, according to Lasky and Williams (2005), the peripheral auditory system is capable of adult-like responses in terms of sensitivity and frequency resolution by about 30 weeks gestational age, with auditory threshold levels primarily adult-like by 35 weeks gestational age (Standley, 2003a, 2003b). By term, newborn sensitivity and frequency resolution is nearly identical to adult levels. However, due to small outer ear canals and immaturity in the middle ear, newborns hear higher frequencies more accurately and attenuate lower frequencies.

Given that the fetus can hear and respond to sound in utero, it is important to consider the auditory sounds present during prenatal development. First and foremost, the mother’s voice is among the most prominent sounds that the fetus hears. The fetus is also exposed to maternal and environmental sounds (Standley, 2003a, 2003b); especially those in the lower frequency range (Lasky & Williams, 2005). In fact, it is now known that the mother’s tissue actually filters the higher frequencies (Lasky & Williams, 2005). Researchers theorize that this emphasis on lower sound frequencies actually helps focus the fetus’ hearing in the area of speech frequencies (Standley, 2003a, 2003b).

Functional development of the auditory system has been studied in both preterm and full term infants, with the instance of congenital hearing loss increasing as gestational age decreases (i.e., the sickest, most preterm newborns have 100 times greater chance of congenital hearing loss) (Lasky & Williams, 2005). This hearing loss has been attributed to exposure to ototoxic drugs, as well as prolonged auditory stimulation (Wroblewska-Seniuk et al., 2005; Zahr & de Traversay, 1995). In fact, neonatal intensive care units can contain excessive amounts of noise and infants are routinely exposed to decibel levels between 55 and 75 dB. Moreover, intermittent noises can reach 100 dB, a sound level comparable to that of a power mower (Lasky & Williams, 2005). Since auditory functioning begins around the 20th week and is adult-like at the 35th week gestational age, premature infants are capable of responding to sounds in their environment. In fact, because of increased oxygen levels used in treating premature infants, premature infants actually display increased auditory sensitivity. As a result, many neonatal intensive care units have begun to reduce environmental stimulation to premature infants (Lasky & Williams, 2005). Furthermore, in order to reduce the negative stressful effects of continuous noise, researchers have called for hospitals to control and/or mask extraneous noise in all areas of patient care (Stouffer & Shirk, 2003). Music, through its unique acoustic properties, has the ability to serve as a masking agent for much of the routine ambient noise in both neonatal intensive care units and general hospital environments (Standley & Whipple, 2003).

### Medical music therapy for preterm infants

Not only can music be used to mask ambient sounds in neonatal intensive care units, it can also be applied therapeutically “to soothe and provide exposure to complex auditory stimuli that promote appropriate neurological stimulation” (Keith, Russell, & Weaver, 2009; Standley & Whipple, 2003, p.21). Before discussing the application of music therapy, it is important to note that music is “acoustically different from all other sound” (Standley, 2003a, 2003b, p. 47) because it is both “sound and silence expressively organized in time” (Madsen & Madsen, 1997, p. 24). Noise, on the other hand, has no fixed pitch (Standley, 2003a, 2003b). Music and noise are processed differently by the brain; noise is irregular and unanticipated, which creates stress. Conversely, music is organized and predictable, functioning to soothe (Standley & Whipple, 2003). Therefore, the unique properties of music can be applied therapeutically by trained music therapists. Music therapy involves the use of evidenced-based music techniques by a trained, credentialed music therapist to achieve behavioral change in an individual (Standley, 2003a, 2003b). Music therapy applications have been shown to be effective in medical settings for a wide range of diagnoses and ages (Standley, 2000). When looking specifically at premature infants, music protocols have been shown to produce the following outcomes: (a) increased oxygen levels, (b) increased sleep time, (c) improved behavior states, (d) decreased hospital stay, (e) reinforced non-nutritive sucking and (f) decreased initial weight loss and improved weight gain (Abromeit, 2003; Keith et

al., 2009; Kemper & Danhauer, 2005; Moore, Gladstone, & Standley, 1994; Standley & Moore, 1995).

### Music therapy protocols

To ensure developmentally appropriate treatment, music therapy protocols have been developed based on gestational age and stage of development. During the survival/pacification stage, which lasts until the infant has reached approximately 2.5 lbs and shows signs of neurological development, protocols primarily consist of non-contact music listening. Infants are exposed to recorded lullabies and their mother's voice during this stage (Standley, 2003a, 2003b). Volume levels are carefully controlled (below 70 dB on scale C) and lullabies are used to promote language development (Standley, 2003a, 2003b). According to Abromeit (2003), studies have been conducted with infants as young as 24 weeks gestational age, with results showing that music applications led to positive health benefits for preterm infants. For example, Caine (1992) found evidence that playing recorded music for infants in the neonatal intensive care unit increased weight gain and shortened lengths of stay. Moreover, a follow-up study of the infants that participated in Caine's music listening study indicated that infants who participated in the music intervention were rated as calmer by their mothers at 6 months of age compared to infants without music intervention (Standley, 1991). Cevasco (2008) also found positive effects for preterm infants who listened to recordings of their mothers' voices; these infants were discharged on average 2 days sooner than infants who did not hear recordings of their mothers' voices. Additionally, Cevasco and Grant (2005) stated that music has been shown to be effective in stabilizing oxygen saturation levels, with no negative effects of apnea/bradycardia noted. Music has even been shown to be more effective than the mother's voice for oxygen stabilization.

Following the survival/pacification stage, infants move into the cautious stimulation phase. During this period, which begins at approximately 30 weeks gestational age, music is used to increase tolerance to stimulation. Several techniques are used during this phase, including music combined with kangaroo care and multimodal stimulation. The first intervention, music combined with kangaroo care (skin to skin contact), has been shown to have positive effects. In fact, research suggests that music combined with kangaroo care can increase occurrences of quiet sleep, decrease crying and significantly lower maternal anxiety (Lai et al., 2006). Multimodal stimulation, which is appropriate for infants starting between 30 and 32 weeks gestational age, involves a musically adapted version of the White-Traut and Tubeszewski (1986) Auditory, Tactile, Vestibular, Visual (ATVV) protocol for reciprocal, multimodal stimulation of premature infants (Standley, 1998). In multimodal stimulation, music paired with stroking, rocking and eye-to-eye contact is used to soothe infants and increase stimulation tolerance (Whipple, 2005). Multimodal stimulation is not appropriate for infants younger than 30 weeks due to hypersensitivity in infants of younger gestational ages. Even with infants of older gestational age, behavioral cues like hiccoughs, tongue protrusion and finger splays must be carefully monitored during treatment in order to avoid overstimulation (Standley, 2003a, 2003b).

In a study of 40 preterm infants who were at least 32 weeks gestational age, Standley (1998) found that female infants receiving multimodal stimulation were discharged significantly sooner (an average of 11.9 days sooner than the no contact control group) and that male infants were discharged 1.5 days sooner. All infants also increased their stimulation tolerance, but again females showed larger gains than males. Whipple (2000) also found that parents who had been taught to use the multimodal procedure increased

their time in the NICU. Additionally, interactions between the premature infants and their parents showed significantly fewer infant stress behaviors and significantly more appropriate parent behaviors (Whipple, 2000).

Preterm infants transition to interactive learning at approximately 32 weeks gestational age (Standley, 2003a, 2003b). During this stage, there are several issues that music therapists can address including inconsolable crying, procedural pain and feeding. Keith et al. (2009) examined the use of developmentally appropriate music interventions to address the issue of inconsolable crying. Infants who have been removed from ventilation but are still unable to receive nutrition orally often demonstrate episodes of inconsolable crying; these episodes can be stressful for infants and parents and can be time consuming in terms of the amount of care provided by the nursing staff. Therefore, Keith et al. (2009) applied developmentally appropriate music listening interventions with infants between the ages of 32 and 40 weeks gestational age in order to determine the effectiveness of music in combating inconsolable crying. Results indicated a significant reduction in the frequency and duration of episodes of inconsolable crying. Results also noted improved physiological measures including heart rate, respiration rate, oxygen saturation, and mean arterial pressure.

Cignacco et al. (2005) conducted a systematic literature review of the efficacy of non-pharmacological interventions in the management of procedural pain in both preterm and term infants. Included in this review was an investigation of the use of music to address procedural pain, specifically in the case of heel sticks. The authors concluded that music invariably produced a positive effect on pain responses. This appeared to be especially true when music was combined with non-nutritive sucking or applied with preterm infants from the ages of 30–31 weeks gestational age or older.

In addition to addressing inconsolable crying and procedural pain, it is important to promote the transition from gavage feeding to nipple feeding during the interactive learning stage. In fact, nipple feeding is one of the criteria that must be met for discharge from most neonatal intensive care units (Kelly, 2006). However, the transition to nipple feeding can be difficult for many preterm infants. In fact, preterm infants often show immature sucking patterns and develop nipple aversion or aversion to oral feeding (Standley, 2003a, 2003b). To encourage nipple acceptance and healthy sucking patterns, music can be used contingently to reinforce non-nutritive sucking (NNS). Non-nutritive sucking has been shown to have a variety of benefits, including promoting neurological development, assisting oral feeding and pacification (Standley & Whipple, 2003). Research suggests that NNS increases oxygen saturation levels when used both prior to and after feeding. Infants who display NNS also have fewer bradycardia and oxygen desaturation episodes. Additionally, NNS led to less behavioral distress, fewer tube feedings and earlier readiness for nipple feeding (Cevasco & Grant, 2005).

Because music can be an effective reinforcer for learning, the application of contingent music to facilitate NNS can be incorporated beginning at approximately 34 weeks gestational age (Standley, 2003a, 2003b). The PAL, or pacifier activated lullaby, provides infants with 10–15 min opportunities one to two times per day to contingently activate music by sucking on a pacifier connected to a pressure transducer and timer (i.e., when the infant sucks in accordance with preset settings, the music turns on). Contingent responses can be altered based on sucking pressure, time intervals and/or desired sucking bursts. Research suggests the application of contingent music is effective in teaching NNS, with most infants capable of attaining 10–15 min of NNS and improving feeding within 1–2 days (Standley, 2003a, 2003b).

Once an infant has successfully completed multimodal stimulation during the cautious stimulation phase and mastered NNS



via PAL during the interactive learning stage, infant stimulation may occur. This procedure involves the use of physical contact (snuggling/cuddling) and stimulating music to facilitate non-crying vocalizations, eye contact and promote language. Additionally, recorded music can be placed in different locations around the infant to encourage head turning to auditory stimuli. Individuals may also provide quiet input (i.e. signing) from a variety of positions in order to stimulate visual focus of attention (Standley, 2003a, 2003b).

A final music therapy technique incorporated into the treatment of premature infants is parental counseling. Parents often experience a wide range of stressors connected with the birth of a premature infant, including financial stressors and care giving stressors (Bachman & Lind, 1997). In a study conducted by Whipple (2000), parents were given multimodal stimulation training as a part of the care giving procedures provided by the NICU staff. Whipple (2000) addressed proper technique, interaction time limits, infant preference for mother's voice (Standley & Madsen, 1990), acceptance of eye-to-eye contact as a sign of contentment/increasing neurological maturity, and the importance of a calming environment and signs of overstimulation (Whipple, 2000). Results indicated that infants of parents who utilized multimodal stimulation showed significantly fewer stress behaviors, displayed greater average daily weight gain, and shorter hospital stays. The parents of these infants also displayed more appropriate interactions and increased time spent in visitation. Therefore parental counseling in music therapy with premature infants strives to promote bonding and goal-oriented interactions, to teach developmental skills and awareness of developmental delays and to provide emotional support for parents when dealing with infant related issues.

## Summary

Music therapy techniques have been demonstrated to be effective across a wide range of medical and developmental issues for premature infants. Due to the unique qualities of music, a trained, board-certified (MT-BC) music therapist with NICU training (NICU-MT) can combine classical music training and a functional knowledge of music with "principles of psychology, education, medicine and rehabilitation" (Standley, 2003a, 2003b, p. 45) to address a wide range of needs with premature infants and their families. Through music therapy techniques like: music listening, music combined with kangaroo care, multimodal stimulation, PAL and parental training, considerable medical and developmental benefits can improve the quality of care in the NICU.

## References

Abromeit, D. H. (2003). The newborn individualized developmental care and assessment program (NIDCAP) as a model for clinical music therapy interventions with premature infants. *Music Therapy Perspectives*, 21, 60–68.

Bachman, D. H., & Lind, R. F. (1997). Perinatal social work and the family of the newborn intensive care infant. *Social Work in Health Care*, 24(3–4), 21–37.

Caine, J. (1992). The effects of music on the selected stress behaviors, weight, caloric and formula intake, and length of hospital stay of premature and low birth weight neonates in a newborn intensive care unit. *Journal of Music Therapy*, 28, 180–192.

Cevasco, A. M. (2008). The effects of mothers' singing on full-term and preterm infants and maternal emotional responses. *Journal of Music Therapy*, 45, 273–306.

Cevasco, A. M., & Grant, R. E. (2005). Effects of the pacifier activated lullaby on weight gain of preterm infants. *Journal of Music Therapy*, 42, 123–139.

Cignacco, E., Hamers, J., Stoffel, L., van Lingen, R., Gessler, P., McDougall, J., & Nelle, M. (2005). The efficacy of non-pharmacological interventions in the management of procedural pain in preterm and term neonates: A systematic literature review. *European Journal of Pain*, 11, 139–152.

Gibson, A., Carney, S., & Wales, J. K. (2006). Growth and the premature baby. *Hormone Research*, 65(Suppl. 3), 75–81.

Hack, M. (2007). Survival and neurodevelopmental outcomes of preterm infants. *Journal of Pediatric Gastroenterology and Nutrition*, 45, S141–S142.

Hepper, P. G., & Shahidullah, B. S. (1994). Development of fetal hearing. *Archives of Disease in Childhood*, 71, F81–F87.

Hoekstra, R. E., Ferrara, T. B., Couser, R. J., Payne, N. R., & Connett, J. E. (2004). Survival and long term neurodevelopmental outcome of extremely premature infants born at 23–26 weeks' gestational age at a tertiary center. *Pediatrics*, 113, e1–e6. Retrieved October 15, 2008 from <http://pediatrics.aappublications.org/cgi/content/abstract/113/1/e1>

Keith, D. R., Russell, K., & Weaver, B. S. (2009). The effects of music listening on inconsolable crying in premature infants. *Journal of Music Therapy*, 46, 191–203.

Kelly, M. M. (2006). Primary care issues for the healthy premature infant. *Journal of Pediatric Health Care*, 20, 293–299.

Kemper, K. J., & Danhauer, S. C. (2005). Music as therapy. *Southern Medical Journal*, 98, 282–288.

Lai, H. L., Chen, C. J., Peng, T. C., Chang, F. M., Hsieh, M. L., Huang, H. Y., & Chang, S. C. (2006). Randomized controlled trial of music during kangaroo care on maternal state anxiety and preterm infants' responses. *International Journal of Nursing Studies*, 43, 139–146.

Lasky, R. E., & Williams, A. L. (2005). The development of the auditory system from conception to term. *NeoReviews*, e141–e152. Retrieved October 15, 2008 from Error! Hyperlink reference not valid. [full/neoreviews/6/3/e141](http://neoreviews/6/3/e141).

Madsen, C. K., & Madsen, C. H. (1997). *Experimental research in music* (3rd ed.). Raleigh, NC: Contemporary Publishing Company of Raleigh, Inc.

Moore, R., Gladstone, I., & Standley, J. (1994, November). *Effects of music, maternal voice, intrauterine sounds and white noise on the oxygen saturation levels of premature infants*. Unpublished paper presented at the National Conference, National Association for Music therapy, Orlando, FL.

Standley, J. M. (1991). Long-term benefits of music intervention in the newborn intensive care unit: A pilot study. *Journal of the International Association of Music for the Handicapped*, 6, 12–23.

Standley, J. M. (1998). The effect of music and multimodal stimulation on responses of premature infants in neonatal intensive care. *Pediatric Nursing*, 24, 532–538.

Standley, J. M. (2000). Music research in medical/dental treatment: An update of a prior meta-analysis. In American Music Therapy Association (Ed.), *Effectiveness of music therapy procedures: Documentation of research and clinical practice* (3rd ed., pp. 1–60). Silver Spring, MD: American Music Therapy Association.

Standley, J. M. (2003). *Music therapy with premature infants: Research and developmental interventions*. Silver Spring, MD: American Music Therapy Association.

Standley, J. (2003). The effect of contingent music to increase non-nutritive sucking of premature infants. *Pediatric Nursing*, 26(493–495), 498–499.

Standley, J., & Madsen, C. (1990). Comparison of infant preferences and responses to auditory stimuli: Music, mother and other female voice. *Journal of Music Therapy*, 27, 54–97.

Standley, J. M., & Moore, R. S. (1995). Therapeutic effects of music and mother's voice on premature infants. *Pediatric Nursing*, 21(509–512), 574.

Standley, J. M., & Whipple, J. (2003). Music therapy for preterm infants in the neonatal intensive care unit: Health and developmental benefits. In S. L. Robb (Ed.), *Music therapy in pediatric healthcare: Research and evidenced-based practice* (pp. 19–30). Silver Spring, MD: American Music Therapy Association.

Stouffer, J. W., & Shirk, B. (2003). Critical care: Applications of music for children on mechanical ventilation. In S. L. Robb (Ed.), *Music Therapy in pediatric healthcare: Research and evidenced-based practice* (pp. 49–80). Silver Spring, MD: American Music Therapy Association.

Whipple, J. (2000). The effect of parent training in music and multimodal stimulation on parent–neonate interactions in the neonatal intensive care unit. *Journal of Music Therapy*, 37, 250–268.

Whipple, J. (2005). Music and multimodal stimulation as developmental intervention in neonatal intensive care. *Music Therapy Perspectives*, 25, 100–110.

White-Traut, R. C., & Tubeszewski, K. A. (1986). Multimodal stimulation of the premature infant. *Journal of Pediatric Nursing*, 1, 1–95.

Wilson-Costello, D., Friedman, H., Minich, N., Fanaroff, A. A., & Hack, M. (2005). Improved survival rates with increased neurodevelopmental disability for extremely low birth weight infants in the 1990s. *Pediatrics*, 115, 997–1003.

Wroblewska-Seniuk, K., Chojnacka, K., Pucher, B., Szczapa, J., Gadzinowski, J., & Gregorowski, M. (2005). The results of newborn hearing screening by means of transient evoked otoacoustic emissions. *International Journal of Pediatric Otorhinolaryngology*, 69, 1351–1357.

Zahr, L. K., & de Traversay, J. (1995). Premature infant responses to noise reduction by earmuffs: Effects on behavioral and physiological measures. *Journal of Perinatology*, 15, 448–455.